


London

Published 1985


Source map scale - 1:25,000

These maps were produced by the Russian military during the Cold War between 1950 and 1997, and cover 103 towns and cities throughout the U.K. The maps are produced at 1:25,000, 1:10,000 and 1:5,000 scale, and show detailed land use, with colour-coded areas for development, green areas, and non-developed areas. Buildings are coloured black and important building uses (such as hospitals, post offices, factories etc.) are numbered, with a numbered key describing their use. They were produced by the Russians for the benefit of navigation, as well as strategic military sites and transport hubs, for use if they were to have invaded the U.K. The detailed information provided indicates that the areas were surveyed using land-based personnel, on the ground, in the cities that are mapped.

Map Name(s) and Date(s)



Russian Map - Slice A



Order Details

Order Number: 186657810_1_1
Customer Ref: 13065 FD
National Grid Reference: 527710, 184060
Slice: A
Site Area (Ha): 0.01
Search Buffer (m): 1000

Site Details

38, Meadowbank, LONDON, NW3 3AY



Ordnance Survey Plan

Published 1991 - 1996


Source map scale - 1:10,000

The historical maps shown were reproduced from maps predominantly held at the scale adopted for England, Wales and Scotland in the 1840's. In 1854 the 1:2,500 scale was adopted for mapping urban areas; these maps were used to update the 1:10,560 maps. The published date given therefore is often some years later than the surveyed date. Before 1938, all OS maps were based on the Cassini Projection, with independent surveys of a single county or group of counties, giving rise to significant inaccuracies in outlying areas. In the late 1940's, a Provisional Edition was produced, which updated the 1:10,560 mapping from a number of sources. The maps appear unfinished - with all military camps and other strategic sites removed. These maps were initially overprinted with the National Grid. In 1970, the first 1:10,000 maps were produced using the Transverse Mercator Projection. The revision process continued until recently, with new editions appearing every 10 years or so for urban areas.

Map Name(s) and Date(s)

TQ28NE
1996
1:10,000
TQ28SE
1991
1:10,000

Historical Map - Slice A



Order Details

Order Number: 186657810_1_1
Customer Ref: 13065 FD
National Grid Reference: 527710, 184060
Slice: A
Site Area (Ha): 0.01
Search Buffer (m): 1000

Site Details

38, Meadowbank, LONDON, NW3 3AY

10k Raster Mapping

Published 1999


Source map scale - 1:10,000

The historical maps shown were produced from the Ordnance Survey's 1:10,000 colour raster mapping. These maps are derived from Landplan which replaced the old 1:10,000 maps originally published in 1970. The data is highly detailed showing buildings, fences and field boundaries as well as all roads, tracks and paths. Road names are also included together with the relevant road number and classification. Boundary information depiction includes county, unitary authority, district, civil parish and constituency.

Map Name(s) and Date(s)

TQ28NE	1999	1:10,000
TQ28SE	1999	1:10,000

Historical Map - Slice A




Order Details

Order Number: 186657810_1_1
Customer Ref: 13065 FD
National Grid Reference: 527710, 184060
Slice: A
Site Area (Ha): 0.01
Search Buffer (m): 1000

Site Details

38, Meadowbank, LONDON, NW3 3AY



10k Raster Mapping

Published 2006

Source map scale - 1:10,000


The historical maps shown were produced from the Ordnance Survey's 1:10,000 colour raster mapping. These maps are derived from Landplan which replaced the old 1:10,000 maps originally published in 1970. The data is highly detailed showing buildings, fences and field boundaries as well as all roads, tracks and paths. Road names are also included together with the relevant road number and classification. Boundary information depiction includes county, unitary authority, district, civil parish and constituency.

Map Name(s) and Date(s)

TQ28NE
2006
1:10,000

TQ28SE
2006
1:10,000

Historical Map - Slice A




Order Details

Order Number: 186657810_1_1
Customer Ref: 13065 FD
National Grid Reference: 527710, 184060
Slice: A
Site Area (Ha): 0.01
Search Buffer (m): 1000

Site Details

38, Meadowbank, LONDON, NW3 3AY





CampbellReith
consulting engineers

VectorMap Local

Published 2018


Source map scale - 1:10,000

VectorMap Local (Raster) is Ordnance Survey's highest detailed 'backdrop' mapping product. These maps are produced from OS's VectorMap Local, a simple vector dataset at a nominal scale of 1:10,000, covering the whole of Great Britain, that has been designed for creating graphical mapping. OS VectorMap Local is derived from large-scale information surveyed at 1:1250 scale (covering major towns and cities), 1:2500 scale (smaller towns, villages and developed rural areas), and 1:10 000 scale (mountain, moorland and river estuary areas).

Map Name(s) and Date(s)

TQ28NE
2018
Variable
TQ28SE
2018
Variable

Historical Map - Slice A



Order Details

Order Number: 186657810_1_1
Customer Ref: 13065 FD
National Grid Reference: 527710, 184060
Slice: A
Site Area (Ha): 0.01
Search Buffer (m): 1000

Site Details



38, Meadowbank, LONDON, NW3 3AY

Landmark
INFORMATION GROUP



Tel: 0844 844 9952
Fax: 0844 844 9951
Web: www.envirocheck.co.uk

Geology 1:10,000 Maps Legends

Artificial Ground and Landslip

Map Colour	Lex Code	Rock Name	Rock Type	Min and Max Age
	WGR	Worked Ground (Undivided)	Unknown/Unclassified Entry	Holocene - Holocene
	MGR	Made Ground (Undivided)	Unknown/Unclassified Entry	Holocene - Holocene

Bedrock and Faults

Map Colour	Lex Code	Rock Name	Rock Type	Min and Max Age
	LC	London Clay Formation	Clay	Eocene - Eocene
	CLGB	Claygate Member	Sandstone	Eocene - Eocene

Geology 1:10,000 Maps

This report contains geological map extracts taken from the BGS Digital Geological map of Great Britain at 1:10,000 scale and is designed for users carrying out preliminary site assessments who require geological maps for the area around a site. This mapping may be more up to date than previously published paper maps.


The various geological layers - artificial and landslip deposits, superficial geology and solid (bedrock) geology are displayed in separate maps, but superimposed on the final 'Combined Surface Geology' map. All map legends feature on this page.

Please Note: Not all of the layers have complete nationwide coverage, so availability of data for relevant map sheets is indicated below.

Geology 1:10,000 Maps Coverage

Map ID:	2	Map ID:	1
Map Name:	TQ28NE	Map Name:	TQ28SE
Map Date:	1999	Map Date:	1999
Bedrock Geology:	Available	Bedrock Geology:	Available
Superficial Geology:	Available	Superficial Geology:	Available
Artificial Geology:	Available	Artificial Geology:	Available
Faults:	Not Available	Faults:	Not Available
Landslip:	Not Available	Landslip:	Not Available
Rock Segments:	Not Available	Rock Segments:	Not Available

Geology 1:10,000 Maps - Slice A



Order Details

Order Number:	186657810_1_1
Customer Ref:	13065 FD
National Grid Reference:	527710, 184060
Slice:	A
Site Area (Ha):	0.01
Search Buffer (m):	1000

Site Details

38, Meadowbank, LONDON, NW3 3AY

Artificial Ground and Landslip


Artificial ground is a term used by BGS for those areas where the ground surface has been significantly modified by human activity. Information about previously developed ground is especially important, as it is often associated with potentially contaminated material, unpredictable engineering conditions and unstable ground.

Artificial ground includes:

- Made ground - man-made deposits such as embankments and spoil heaps on the natural ground surface.
- Worked ground - areas where the ground has been cut away such as quarries and road cuttings.
- Infilled ground - areas where the ground has been cut away then wholly or partially backfilled.
- Landscaped ground - areas where the surface has been reshaped.
- Disturbed ground - areas of ill-defined shallow or near surface mineral workings where it is impracticable to map made and worked ground separately.

Mass movement (landslip) deposits on BGS geological maps are primarily superficial deposits that have moved down slope under gravity to form landslips. These affect bedrock, other superficial deposits and artificial ground. The dataset also includes founded strata, where the ground has collapsed due to subsidence.

Artificial Ground and Landslip Map - Slice A




Order Details

Order Number: 186657810_1_1
 Customer Ref: 13065 FD
 National Grid Reference: 527710, 184060
 Slice: A
 Site Area (Ha): 0.01
 Search Buffer (m): 1000

Site Details

38, Meadowbank, LONDON, NW3 3AY




Superficial Geology

BGS 1:10,000 Superficial Deposits are the youngest geological deposits formed during the most recent period of geological time, which extends back about 1.8 million years from the present.

They rest on older deposits or rocks referred to as Bedrock. This dataset contains Superficial deposits that are of natural origin and 'in place'. Other superficial strata may be held in the Mass Movement dataset where they have been moved, or in the Artificial Ground dataset where they are of man-made origin.

Most of these Superficial deposits are unconsolidated sediments such as gravel, sand, silt and clay, and onshore they form relatively thin, often discontinuous patches or larger spreads.

Superficial Geology Map - Slice A




Order Details

Order Number: 186657810_1_1
 Customer Ref: 13065 FD
 National Grid Reference: 527710, 184060
 Slice: A
 Site Area (Ha): 0.01
 Search Buffer (m): 1000

Site Details

38, Meadowbank, LONDON, NW3 3AY



Bedrock and Faults


Bedrock geology is a term used for the main mass of rocks forming the Earth and are present everywhere, whether exposed at the surface in outcrops or concealed beneath superficial deposits or water.

The bedrock has formed over vast lengths of geological time ranging from ancient and highly altered rocks of the Proterozoic, some 2500 million years ago, or older, up to the relatively young Pliocene, 1.8 million years ago.

The bedrock geology includes many lithologies, often classified into three types based on origin: igneous, metamorphic and sedimentary.

The BGS Faults and Rock Segments dataset includes geological faults and thin beds mapped as lines such as coal seams and mineral veins. These are not restricted by age and could relate to features of any of the 1:10,000 geology datasets.

Bedrock and Faults Map - Slice A



Order Details

Order Number: 186657810_1_1
Customer Ref: 13065 FD
National Grid Reference: 527710, 184060
Slice: A
Site Area (Ha): 0.01
Search Buffer (m): 1000

Site Details

38, Meadowbank, LONDON, NW3 3AY

