


PARTICLE SIZE DISTRIBUTION TEST

In accordance with clauses 9.2 of BS1377:Part 2:1990


Trial Pit: **TP1** Sample Ref: **-** Sample Type: **D** Depth (m): **1.60**



CLAY	fine	medium	coarse	fine	medium	coarse	fine	medium	coarse	COBBLES
	-	-	-	3%	27%	10%	9%	25%	23%	
	SILT			SAND			GRAVEL			
4%				39%			57%			0%

Test Sieve (mm)	Percent Passing (%)	Particle Diameter (mm)	Percent Passing (%)	Coefficients	
125.0	100			D ₁₀ (mm)	0.246
75.0	100			D ₁₅ (mm)	0.315
63.0	100			D ₃₀ (mm)	0.550
37.5	100			D ₅₀ (mm)	5.104
20.0	77			D ₆₀ (mm)	10.393
10.0	59			D ₈₅ (mm)	24.888
6.30	52			D ₉₀ (mm)	28.532
3.35	46			C _U	42
2.00	43			C _C	0.12
1.18	41			Sedimentation sample was not pre-treated Soil Description: Orangish brown mottle white and dark grey very sandy slightly clayey GRAVEL	
0.600	33				
0.425	21				
0.212	7				
0.150	5				
0.063	4				

Key: C_U = Uniformity coefficient. C_C = Coefficient of curvature as defined in BS EN ISO 14688-2



STRUCTURAL SOILS
18 Frogmore Road
Hemel Hempstead
Hertfordshire
HP3 9RT


Compiled By		Date
SHARON CAIRNS		19/03/20
Contract		Contract Ref:
Grange Central St Martins		584233



PARTICLE SIZE DISTRIBUTION TEST

In accordance with clauses 9.2 of BS1377:Part 2:1990


Trial Pit: **TP4** Sample Ref: **-** Sample Type: **D** Depth (m): **1.00**



CLAY	fine	medium	coarse	fine	medium	coarse	fine	medium	coarse	COBBLES
	-	-	-	2%	19%	7%	11%	29%	27%	
	SILT			SAND			GRAVEL			
5%				28%			67%			0%

Test Sieve (mm)	Percent Passing (%)	Particle Diameter (mm)	Percent Passing (%)	Coefficients	
125.0	100			D ₁₀ (mm)	0.256
75.0	100			D ₁₅ (mm)	0.352
63.0	100			D ₃₀ (mm)	1.180
37.5	98			D ₅₀ (mm)	8.573
20.0	73			D ₆₀ (mm)	12.746
10.0	53			D ₈₅ (mm)	27.044
6.30	44			D ₉₀ (mm)	30.667
3.35	37			C _U	50
2.00	33			C _C	0.43
1.18	30			Sedimentation sample was not pre-treated Soil Description: Brown mottled dark grey and light brown very sandy slightly clayey GRAVEL	
0.600	26				
0.425	18				
0.212	7				
0.150	6				
0.063	5				

Key: C_U = Uniformity coefficient. C_C = Coefficient of curvature as defined in BS EN ISO 14688-2



STRUCTURAL SOILS
18 Frogmore Road
Hemel Hempstead
Hertfordshire
HP3 9RT


Compiled By		Date
<i>SC</i>		19/03/20
Contract		Contract Ref:
Grange Central St Martins		584233



PARTICLE SIZE DISTRIBUTION TEST

In accordance with clauses 9.2 of BS1377:Part 2:1990


Trial Pit: **TP5** Sample Ref: **-** Sample Type: **D** Depth (m): **0.70**



CLAY	fine	medium	coarse	fine	medium	coarse	fine	medium	coarse	COBBLES
	-	-	-	2%	24%	10%	13%	30%	18%	
	SILT			SAND			GRAVEL			
4%				35%			61%			0%

Test Sieve (mm)	Percent Passing (%)	Particle Diameter (mm)	Percent Passing (%)	Coefficients	
125.0	100			D ₁₀ (mm)	0.255
75.0	100			D ₁₅ (mm)	0.322
63.0	100			D ₃₀ (mm)	0.661
37.5	100			D ₅₀ (mm)	5.380
20.0	82			D ₆₀ (mm)	9.117
10.0	62			D ₈₅ (mm)	22.209
6.30	52			D ₉₀ (mm)	26.446
3.35	44			C _U	36
2.00	39			C _C	0.19
1.18	36			Sedimentation sample was not pre-treated Soil Description: Orangish brown mottled dark grey and white very sandy slightly clayey GRAVEL	
0.600	29				
0.425	21				
0.212	6				
0.150	5				
0.063	4				

Key: C_U = Uniformity coefficient. C_C = Coefficient of curvature as defined in BS EN ISO 14688-2



STRUCTURAL SOILS
18 Frogmore Road
Hemel Hempstead
Hertfordshire
HP3 9RT


Compiled By		Date
<i>SC</i>		19/03/20
Contract		Contract Ref:
Grange Central St Martins		584233



PARTICLE SIZE DISTRIBUTION TEST

In accordance with clauses 9.2 of BS1377:Part 2:1990


Trial Pit: **TP7** Sample Ref: **-** Sample Type: **D** Depth (m): **0.50**



CLAY	fine	medium	coarse	fine	medium	coarse	fine	medium	coarse	COBBLES
	-	-	-	1%	21%	6%	11%	30%	28%	
	SILT			SAND			GRAVEL			
3%				28%			69%			0%

Test Sieve (mm)	Percent Passing (%)	Particle Diameter (mm)	Percent Passing (%)	Coefficients	
125.0	100			D ₁₀ (mm)	0.286
75.0	100			D ₁₅ (mm)	0.366
63.0	100			D ₃₀ (mm)	1.536
37.5	96			D ₅₀ (mm)	9.117
20.0	72			D ₆₀ (mm)	13.195
10.0	52			D ₈₅ (mm)	28.113
6.3	42			D ₉₀ (mm)	32.047
3.35	36			C _U	46
2.0	31			C _C	0.63
1.18	29			Sedimentation sample was not pre-treated Soil Description: Orangish brown mottled dark grey, brown and white very sandy slightly clayey GRAVEL	
0.600	25				
0.425	18				
0.212	4				
0.150	3				
0.063	3				

Key: C_U = Uniformity coefficient. C_C = Coefficient of curvature as defined in BS EN ISO 14688-2




STRUCTURAL SOILS
18 Frogmore Road
Hemel Hempstead
Hertfordshire
HP3 9RT

Compiled By		Date
SHARON CAIRNS		19/03/20
Contract		Contract Ref:
Grange Central St Martins		584233



In accordance with clauses 9.2 of BS1377:Part 2:1990

Depth (m): **1.90**



CLAY	fine	medium	coarse	fine	medium	coarse	fine	medium	coarse	COBBLES
	-	-	-	3%	29%	3%	5%	28%	30%	
	SILT			SAND			GRAVEL			
2%				35%			63%			0%


Test Sieve (mm)	Percent Passing (%)
125.0	100
75.0	100
63.0	100
37.5	94
20.0	70
10.0	51
6.30	42
3.35	38
2.00	37
1.18	36
0.600	34
0.425	30
0.212	5
0.150	3
0.063	2

Particle Diameter (mm)	Percent Passing (%)
Sedimentation sample was not pre-treated	

Coefficients	
D_{10} (mm)	0.244
D_{15} (mm)	0.280
D_{30} (mm)	0.425
D_{50} (mm)	9.500
D_{60} (mm)	13.887
D_{85} (mm)	29.625
D_{90} (mm)	33.770
C_U	57
C_C	0.05

Orangish brown mottled dark grey and white very sandy slightly clayey GRAVEL

Key: C_u = Uniformity coefficient. C_c = Coefficient of curvature as defined in BS EN ISO 14688-2



Compiled By

Date _____


Contract Ref:

584233



In accordance with clauses 9.2 of BS1377:Part 2:1990

Depth (m): **1.90**



Key: C_U = Uniformity coefficient. C_C = Coefficient of curvature as defined in BS EN ISO 14688-2





APPENDIX I

WM3 ASSESSMENT

Haswaste, developed by Dr. Iain Haslock.

Site Code and Name
TP/WS/BH Depth (m) Envirolab reference

TP12	TP3	TP4						
0.40	1.50	0.50						
20/02615/1	20/01781/1	20/01781/2						

% Moisture	%	2.2	25.3	8.4					
pH (soil)									
pH (leachate)									
Arsenic	mg/kg	5	53	12					
Cadmium	mg/kg	<0.5	<0.5	<0.5					
Copper	mg/kg	8	650	36					
CrVI or Chromium	mg/kg	24	18	68					
Lead	mg/kg	15	4,680	225					
Mercury	mg/kg	<0.17	26.50	0.52					
Nickel	mg/kg	12	23	49					
Selenium	mg/kg	<1	<1	<1					
Zinc	mg/kg	36	209	192					
Barium	mg/kg								
Beryllium	mg/kg								
Vanadium	mg/kg								
Cobalt	mg/kg								
Manganese	mg/kg								
Molybdenum	mg/kg								
Antimony	mg/kg								
Aluminium	mg/kg								
Bismuth	mg/kg								
CrIII	mg/kg								
Iron	mg/kg								
Strontium	mg/kg								
Tellurium	mg/kg								
Thallium	mg/kg								
Titanium	mg/kg								
Tungsten	mg/kg								
Ammoniacal N	mg/kg								
ws Boron	mg/kg								

PAH (Input Total PAH OR individual PAH results)		mg/kg	mg/kg	mg/kg					
Acenaphthene	mg/kg	<0.01	<0.01	<0.01					
Acenaphthylene	mg/kg	<0.01	<0.01	<0.01					
Anthracene	mg/kg	0.03	<0.02	<0.02					
Benzo(a)anthracene	mg/kg	0.08	<0.04	0.10					
Benzo(a)pyrene	mg/kg	0.05	<0.04	0.08					
Benzo(b)fluoranthene	mg/kg	0.07	<0.05	0.11					
Benzo(ghi)perylene	mg/kg	<0.05	<0.05	0.08					
Benzo(k)fluoranthene	mg/kg	<0.07	<0.07	<0.07					
Chrysene	mg/kg	0.07	<0.06	0.11					
Dibenzo(ah)anthracene	mg/kg	<0.04	<0.04	<0.04					
Fluoranthene	mg/kg	0.24	<0.08	0.13					
Fluorene	mg/kg	<0.01	<0.01	<0.01					
Indeno(123cd)pyrene	mg/kg	0.03	0.04	0.07					
Naphthalene	mg/kg	<0.03	<0.03	<0.03					
Phenanthrene	mg/kg	0.16	<0.03	0.05					
Pyrene	mg/kg	0.16	<0.07	0.10					
Coronene	mg/kg	<0.01	0.04	0.03					
Total PAHs (16 or 17)	mg/kg	0.89	0.08	0.86					

TPH									
Petrol	mg/kg								
Diesel	mg/kg								
Lube Oil	mg/kg								
Crude Oil	mg/kg								
White Spirit / Kerosene	mg/kg								
Creosote	mg/kg								
Unknown TPH with ID	mg/kg								
Unknown TPHCWG	mg/kg								
Total Sulphide	mg/kg								
Complex Cyanide	mg/kg								
Free (or Total) Cyanide	mg/kg								
Thiocyanate	mg/kg								
Elemental/Free Sulphur	mg/kg								

Phenols								
Input Total Phenols HPLC OR individual Phenol results.								
Phenol	mg/kg							
Cresols	mg/kg							
Xylenols	mg/kg							
Resourcinol	mg/kg							
Phenols Total by HPLC	mg/kg							

BTEX Input Total BTEX OR individual BTEX results.								
Benzene	mg/kg							
Toluene	mg/kg							
Ethylbenzene	mg/kg							
Xylenes	mg/kg							
Total BTEX	mg/kg							

PCBs (POPs)								
PCBs Total (eq EC7/WHO12)	mg/kg							

[illegible]



Please enter available data in the rows associated with the test (grey) cells. Calculation cells initially display either "0.0000" or "#DIV/0!".
If any calculation cells below state "0.00000", testing has NOT been undertaken that contributes to that Hazardous Property.

Haswaste, developed by Dr. Iain Haslock.

Site Code and Name

TP/WS/BH

Depth (m)

Envirolab reference

TP12	TP3	TP4						
0.40	1.50	0.50						
20/02615/1	20/01781/1	20/01781/2						

POPs Dioxins and Furans Input Total Dioxins and Furans

OR individual Dioxin and Furan results.

2,3,7,8-TeCDD	mg/kg							
1,2,3,7,8-PeCDD	mg/kg							
1,2,3,4,7,8-HxCDD	mg/kg							
1,2,3,6,7,8-HxCDD	mg/kg							
1,2,3,7,8,9-HxCDD	mg/kg							
1,2,3,4,6,7,8-HpCDD	mg/kg							
OCDD	mg/kg							
2,3,7,8-TeCDF	mg/kg							
1,2,3,7,8-PeCDF	mg/kg							
2,3,4,7,8-PeCDF	mg/kg							
1,2,3,4,7,8-HxCDF	mg/kg							
1,2,3,6,7,8-HxCDF	mg/kg							
2,3,4,6,7,8-HxCDF	mg/kg							
1,2,3,7,8,9-HxCDF	mg/kg							
1,2,3,4,6,7,8-HpCDF	mg/kg							
1,2,3,4,7,8,9-HpCDF	mg/kg							
OCDF	mg/kg							
Total Dioxins and Furans	mg/kg							

Some Pesticides (POPs unless otherwise stated)

Aldrin	mg/kg							
α Hexachlorocyclohexane (alpha-HCH) (leave empty if total HCH results used)	mg/kg							
β Hexachlorocyclohexane (beta-HCH) (leave empty if total HCH results used)	mg/kg							
α Cis-Chlordane (alpha) OR Total Chlordane	mg/kg							
δ Hexachlorocyclohexane (delta-HCH) (leave empty if total HCH results used)	mg/kg							
Dieldrin	mg/kg							
Endrin	mg/kg							
γ Hexachlorocyclohexane (gamma-HCH) (lindane) OR Total HCH	mg/kg							
Heptachlor	mg/kg							
Hexachlorobenzene	mg/kg							
o,p'-DDT (leave empty if total DDT results used)	mg/kg							
p,p'-DDT OR Total DDT	mg/kg							
γ Trans-Chlordane (gamma) (leave empty if total Chlordane results used)	mg/kg							
Chlordecone (kepone)	mg/kg							
Pentachlorobenzene	mg/kg							
Mirex	mg/kg							
Toxaphene (camphechlor)	mg/kg							

Tin

Tin (leave empty if Organotin and Tin excl Organotin results used)

Organotin

Dibutyltin; DiBT	mg/kg							
Tributyltin; TriBT	mg/kg							
Triphenyltin; TriPT	mg/kg							
Tetrabutyltin; TeBT	mg/kg							
Tin excluding Organotin								
Tin excl Organotin	mg/kg							



Haswaste, developed by Dr. Iain Haslock.

Site Code and Name

TP/WS/BH

Depth (m)

Envirolab reference

Asbestos in Soil

Asbestos detected in Soil (enter Y or N)

Thresholds

Y

Asbestos % Composition in Soil (Matrix Loose Fibres or Microscopic Identifiable Pieces only)

see "Carc HP7 % Asbestos in Soil (Fibres)" below

%

Carcinogenic HP7 % Asbestos in Soil (fibres or micro pieces)

Please be advised, if the calculation cell is "0.00000" DOES NOT MEAN asbestos testing has been undertaken and the result is zero.

≥0.1%

Asbestos Identifiable Pieces visible with the naked eye detected in the Soil (enter Y or N)

Y

Please enter available data in the rows associated with the test (grey) cells. Calculation cells initially display either "0.0000" or "#DIV/0!".
If any calculation cells below state "0.00000", testing has NOT been undertaken that contributes to that Hazardous Property.

TP12	TP3	TP4						
0.40	1.50	0.50						
20/02615/1	20/01781/1	20/01781/2						

				Y				
--	--	--	--	---	--	--	--	--

If Asbestos in Soil above is "Y", the soil is Hazardous Waste HP5 and HP7

0.00000	0.00000	0.00000	0.00000	0.00000	0.00000	0.00000	0.00000	0.00000

If Asbestos in Soil above is "Y", but Asbestos % above is "<0.1%", the soil is Non Hazardous Waste. You can only use Asbestos % results where loose fibres or micro pieces are only present. You cannot use Asbestos % results when visual identifiable pieces are present.

--	--	--	--	--	--	--	--	--

If visual identifiable pieces of asbestos are present, you cannot use Asbestos % results and the whole soil sample is Hazardous Waste HP5 and HP7 Construction material containing Asbestos 17 06 05. Therefore, if Asbestos in Soil above is "Y", the Asbestos % above is "<0.1%", but the Asbestos Identifiable Pieces visible with the naked eye is "Y", the soil is Hazardous Waste.

Identifiable Pieces are Cement, Fragments, Board, Rope etc. ie anything ACM that is not Loose Fibres.

All visual asbestos pieces need to be removed leaving only fibres (or micro pieces) with an Asbestos % Composition in Soil result of <0.1% for the soil to become non-hazardous waste.

Hazardous Property	Thresholds	Cut Off Value	If cells below turn yellow and the text turns red, the samples should be classified as Hazardous Waste.					
Corrosive HP8	≥5%	<1%	0.00515	0.00781	0.01341	0.00000	0.00000	0.00000
Irritant HP4	≥10%	<1%	0.00153	0.06009	0.00518	0.00000	0.00000	0.00000
Irritant HP4	≥20%	<1%	#VALUE!	#VALUE!	#VALUE!	0.00000	0.00000	0.00000
Specific Target Organ Toxicity HP5	≥1%		0.00000	0.00000	0.00000	0.00000	0.00000	0.00000
Specific Target Organ Toxicity HP5	≥20%		0.00009	0.00001	0.00008	0.00000	0.00000	0.00000
Specific Target Organ Toxicity HP5	≥1%		0.00451	0.00347	0.01196	0.00000	0.00000	0.00000
Specific Target Organ Toxicity HP5	≥10%		#VALUE!	#VALUE!	#VALUE!	0.00000	0.00000	0.00000
Aspiration Toxicity HP5	≥10%		0.00000	0.00000	0.00000	0.00000	0.00000	0.00000
Acute Toxicity HP6	≥0.1%	<0.1%	0.00000	0.00000	0.00000	0.00000	0.00000	0.00000
Acute Toxicity HP6	≥0.25%	<0.1%	#VALUE!	0.00721	0.00150	0.00000	0.00000	0.00000
Acute Toxicity HP6	≥5%	<0.1%	#VALUE!	#VALUE!	#VALUE!	0.00000	0.00000	0.00000
Acute Toxicity HP6	≥25%	<1%	#VALUE!	#VALUE!	#VALUE!	0.00000	0.00000	0.00000
Acute Toxicity HP6	≥0.25%	<0.1%	#VALUE!	0.00198	0.00005	0.00000	0.00000	0.00000
Acute Toxicity HP6	≥2.5%	<0.1%	0.00451	0.00258	0.01196	0.00000	0.00000	0.00000
Acute Toxicity HP6	≥15%	<0.1%	0.00000	0.00000	0.00000	0.00000	0.00000	0.00000
Acute Toxicity HP6	≥55%	<1%	#VALUE!	#VALUE!	#VALUE!	0.00000	0.00000	0.00000
Acute Toxicity HP6	≥0.1%	<0.1%	0.00000	0.00000	0.00000	0.00000	0.00000	0.00000
Acute Toxicity HP6	≥0.5%	<0.1%	#VALUE!	#VALUE!	#VALUE!	0.00000	0.00000	0.00000
Acute Toxicity HP6	≥3.5%	<0.1%	#VALUE!	#VALUE!	#VALUE!	0.00000	0.00000	0.00000
Acute Toxicity HP6	≥22.5%	<1%	0.00472	0.40793	0.03340	0.00000	0.00000	0.00000
Carcinogenic HP7	≥0.1%		0.00451	0.34960	0.02061	0.00000	0.00000	0.00000
Carcinogenic HP7	≥0.1%		0.000000000	0.000000000	0.000000000	0.000000000	0.000000000	0.000000000
Carcinogenic HP7	≥1%		0.00009	0.00001	0.00008	0.00000	0.00000	0.00000
Carcinogenic HP7 Unknown TPH with ID	≥1,000mg/kg		0.00	0.00	0.00	0.00	0.00	0.00
Carcinogenic HP7 b(a)p marker test (Unknown TPH with ID only) Cell only applicable if TPH >1,000mg/kg	≥0.01%		#DIV/0!	#DIV/0!	#DIV/0!	#DIV/0!	#DIV/0!	#DIV/0!
pH Corrosive HP8 pH (soil or leachate)	H8 ≥11.5		0.00	0.00	0.00	0.00	0.00	0.00
pH Corrosive HP8 pH (soil or leachate)	H8 ≤2		0.00	0.00	0.00	0.00	0.00	0.00
Toxic for Reproduction HP10	≥0.3%		0.00237	0.34960	0.02061	0.00000	0.00000	0.00000
Toxic for Reproduction HP10	≥3%		0.00451	0.00258	0.01196	0.00000	0.00000	0.00000
Mutagenic HP11	≥0.1%		0.00451	0.00258	0.01196	0.00000	0.00000	0.00000
Mutagenic HP11 Unknown TPH with ID	≥1,000mg/kg		0.00	0.00	0.00	0.00	0.00	0.00
Mutagenic HP11 b(a)p marker test (Unknown TPH with ID only) Cell only applicable if TPH >1,000mg/kg	≥0.01%		#DIV/0!	#DIV/0!	#DIV/0!	#DIV/0!	#DIV/0!	#DIV/0!
Mutagenic HP11	≥1%		0.00237	0.00347	0.00907	0.00000	0.00000	0.00000
Produces Toxic Gases HP12 Sulphide	≥1,400mg/kg		0.0	0.0	0.0	0.0	0.0	0.0
Produces Toxic Gases HP12 Cyanide	≥1,200mg/kg		0.0	0.0	0.0	0.0	0.0	0.0
Produces Toxic Gases HP12 Thiocyanate	≥2,600mg/kg		0.0	0.0	0.0	0.0	0.0	0.0
HP13 Sensitising	≥10%		0.00451	0.00347	0.01196	0.00000	0.00000	0.00000



Please enter available data in the rows associated with the test (grey) cells. Calculation cells initially display either "0.0000" or "#DIV/0!".
If any calculation cells below state "0.00000", testing has NOT been undertaken that contributes to that Hazardous Property.

Haswaste, developed by Dr. Iain Haslock.

Site Code and Name

TP/WS/BH

Depth (m)

Envirolab reference

TP12	TP3	TP4						
0.40	1.50	0.50						
20/02615/1	20/01781/1	20/01781/2						

Ecotoxic HP14 amended v6	≥25%	<0.1%	#VALUE!	#VALUE!	#VALUE!	0.00000	0.00000	0.00000	0.00000	0.00000	0.00000	0.00000	0.00000	0.00000
Ecotoxic HP14 amended v6	≥25%	<0.1% (except Be, V, Te, Ti, Petrol, Diesel, Crude Oil, Kerosene, White Spirit, Crosole, TPH, TPHCWG, Phenol, Cresols, Xylenols, T-Phenols, CompCN, Thiocyanate, Toluene, Ethylbenzene, Xylene + BTEX 1%).	#VALUE!	#VALUE!	#VALUE!	0.00000	0.00000	0.00000	0.00000	0.00000	0.00000	0.00000	0.00000	0.00000
Ecotoxic HP14 amended v6	≥25%	<0.1% (except Be, V, Te, Ti, Petrol, Diesel, Crude Oil, Kerosene, White Spirit, Crosole, TPH, TPHCWG, Phenol, Cresols, Xylenols, T-Phenols, CompCN, Thiocyanate, Toluene, Ethylbenzene, Xylene + BTEX 1%).	#VALUE!	#VALUE!	#VALUE!	0.00000	0.00000	0.00000	0.00000	0.00000	0.00000	0.00000	0.00000	0.00000
Persistent Organic Pollutant (PCB, PBB or POP Pesticides)	>0.005%		0.00000000	0.00000000	0.00000000	0.00000000	0.00000000	0.00000000	0.00000000	0.00000000	0.00000000	0.00000000	0.00000000	0.00000000
Persistent Organic Pollutant (Total Dioxins+Furans)	>0.0000015%		0.0000000000	0.0000000000	0.0000000000	0.0000000000	0.0000000000	0.0000000000	0.0000000000	0.0000000000	0.0000000000	0.0000000000	0.0000000000	0.0000000000
Persistent Organic Pollutant (Individual Dioxins+Furans)	>0.0000015%		0.0000000000	0.0000000000	0.0000000000	0.0000000000	0.0000000000	0.0000000000	0.0000000000	0.0000000000	0.0000000000	0.0000000000	0.0000000000	0.0000000000



APPENDIX J

RSK SOIL GACS

Generic assessment criteria for human health: residential scenario without home-grown produce

Background

RSK's generic assessment criteria (GAC) were initially prepared following the publication by the Environment Agency (EA) of soil guideline value (SGV) and toxicological (TOX) reports, and associated publications in 2009⁽¹⁾. RSK GAC were updated following the publication of GAC by LQM/CIEH in 2009⁽²⁾. RSK GAC are periodically revised when updated information on toxicological, land use or receptor parameters is published.

Updates to the RSK GAC

In 2014, the publication of Category 4 Screening Levels (C4SL)^(3,4), as part of the Defra-funded research project SP1010, included modifications to certain exposure assumptions documented within EA Science Report SC050221/SR3 (herein after referred to as SR3)⁽⁵⁾ used in the generation of SGVs.

C4SL were published for six substances (cadmium, arsenic, benzene, benzo(a)pyrene, chromium VI and lead) for a sandy loam soil type with 6% soil organic matter, based on a low level of toxicological concern (LLTC; see Section 2.3 of research project report SP1010⁽³⁾). Where a C4SL has been published, the RSK GAC duplicates the C4SL published values using all input parameters within the SP1010 final project report⁽³⁾ and associated appendices⁽⁶⁾, and adopts them as GAC for these six substances.

For all other substances the C4SL exposure modifications relevant for residential without home-grown produce end use have been applied to the current RSK GAC. These include alterations to daily inhalation rates for residential and commercial scenarios, reducing soil adherence factors in children (age classes 1 to 12 only) and reducing exposure frequency for dermal contact outdoors.

The RSK GAC have also been revised with updated toxicology published by LQM/CIEH in 2015⁽⁷⁾ or by the USEPA⁽¹⁴⁾, where a C4SL has not been published.

RSK GAC derivation for metals and organic compounds

Model selection

Soil assessment criteria (SAC) were calculated using the Contaminated Land Exposure Assessment (CLEA) tool v1.071, supporting EA guidance^(5,8,9) and revised exposure scenarios published for the C4SL⁽³⁾. The SAC are also termed GAC.

Conceptual model

In accordance with SR3⁽⁵⁾, the residential without home-grown produce scenario considers risks to a female child between the ages of 0 and 6 years old as the highest risk scenario. In accordance with Box 3.1 of SR3⁽⁵⁾, the pathways considered for production of the SAC in the residential without home-grown produce scenario are

- direct soil and dust ingestion in areas of soft landscaping
- dermal contact with soil and indoor dust

- inhalation of indoor and outdoor dust and vapours.

Figure 1 is a conceptual model illustrating these linkages.

In line with guidance in the EA SGV report for cadmium⁽¹⁾, the RSK GAC for cadmium has been derived based on estimates representative of lifetime exposure. Although young children are generally more likely to have higher exposures to soil contaminants, the renal toxicity of cadmium, and the derivation of the TDI_{oral} and TDI_{inh} , are based on considerations of the kidney burden accumulated over 50 years or so. It is therefore reasonable to consider exposure not just in childhood but averaged over a longer period.

With respect to volatilisation, the CLEA model assumes a simple linear partitioning of a chemical in the soil between the sorbed, dissolved and vapour phase⁽⁹⁾. The upper boundaries of this partitioning are represented by the maximum aqueous solubility and pure saturated vapour concentration of the chemical. The CLEA model estimates saturated soil concentrations where these limits are reached⁽⁹⁾. The CLEA software uses a traffic light system to identify when individual and/or combined assessment criteria exceed the lower of either the aqueous- or vapour-based soil saturation limits. Model output cells are flagged red where the saturated soil concentration has been exceeded and the contribution of the indoor and outdoor vapour pathway to total exposure is greater than 10%. In this case, further consideration of the following is required⁽⁹⁾:

- Free phase contamination may be present.
- Exposure from the vapour pathways will be over-predicted by the model, as in reality the vapour phase concentration will not increase at concentrations above saturation limits
- Where the vapour pathway contribution is greater than 90%, it is unlikely the relevant health criteria value (HCV) will be exceeded at soil concentrations at least a factor of ten higher than the relevant HCV.

Where the vapour pathway is the predominant pathway (contributes greater than 90% of exposure) or the only exposure route considered and the cell is highlighted red (SAC exceeds saturation limit), the risk based on the assumed conceptual model is likely to be negligible as the vapour risk is assumed to be tolerable at maximum possible soil concentrations. In such circumstances, the vapour pathway exposure should be considered based on the presence of free phase or non-aqueous phase liquid sources and the measured concentrations of volatile organic compounds (VOC) in the vapour phase. Screening could be considered based on setting the SAC as the modelled soil saturation limits. However, as stated within the CLEA handbook⁽⁹⁾, this is likely to not be practical in many cases because of the very low saturation limits and, in any case, is highly conservative.

It should also be noted that for mixtures of compounds, free phase may be present where soil (or groundwater) concentrations are well below saturation limits for individual compounds.

Where the vapour pathway is only one of the exposure pathways considered, an additional approach can then be utilised as detailed within Section 4.12 of the CLEA model handbook⁽⁹⁾, which explains how to calculate an effective assessment criterion manually.

SR3⁽⁵⁾ states that, as a general rule of thumb, it is recognised that estimating vapour phase concentrations from dissolved and sorbed phase contamination by petroleum hydrocarbons are at least a factor of ten higher than those likely to be measured on-site. RSK has therefore applied an empirical subsurface to indoor air correction factor of 10 into the CLEA model chemical database for all petroleum hydrocarbon fractions (including BTEX, trimethylbenzenes and the

polycyclic aromatic hydrocarbons (PAH) naphthalene, acenaphthene and acenaphthylene) to reduce this conservatism.

Input selection

The most up-to-date published chemical and toxicological data was obtained from EA Report SC050021/SR7⁽¹⁰⁾, the EA TOX⁽¹¹⁾ reports, the C4SL SP1010 project report and associated appendices^(3,6), the 2015 LQM/CIEH report⁽⁷⁾ or the USEPA IRIS database⁽¹⁴⁾. Where a C4SL has been published, the RSK GAC have duplicated the C4SL published values using all input parameters within the SP1010 final project report⁽³⁾ and associated appendices⁽⁶⁾, and has adopted them as GAC for these six substances. Toxicological and specific chemical parameters for 1,2,4-trimethylbenzene, barium and methyl tertiary-butyl ether (MTBE) were obtained from the CL:AIRE Soil Generic Assessment Criteria report⁽¹¹⁾.

For TPH, aromatic hydrocarbons C₅–C₈ were not modelled, as this range comprises benzene (>EC5-EC7) and toluene (>EC7-EC8), which are modelled separately.

Physical parameters

For the residential without home-grown produce scenario, the CLEA default building is a small, two-storey terrace house with a concrete ground-bearing slab. SR3⁽⁵⁾ notes this residential building type to be the most conservative in terms of potential for vapour intrusion. The building parameters used in the production of the RSK GACs are the default CLEA v1.06 inputs presented in Table 3.3 of SR3⁽³⁾, with a dust loading factor detailed in Section 9.3 of SR3⁽⁵⁾. The parameters for a sandy loam soil type were used in line with Table 4.4 of SR3⁽⁵⁾. This includes a value of 6% for the percentage of soil organic matter (SOM) within the soil. In RSK's experience, this is rather high for many sites. To avoid undertaking site-specific risk assessments for this SOM, RSK has produced an additional set of GAC for SOM of 1% and 2.5% for all substances using the CLEA tool.

Summary of modifications to the default CLEA SR3⁽⁵⁾ input parameters for residential without home-grown produce

In summary, the RSK GAC were produced using the default input parameters for soil properties, the air dispersion model, building properties and the vapour model detailed in SR3⁽⁵⁾. Modifications to the default SR3⁽⁵⁾ exposure scenarios based on the C4SL exposure scenarios⁽³⁾ are presented in Table 2 below.

The final selected GAC are presented by pathway in Table 3 and the combined GAC in Table 4.

Figure 1: Conceptual model for CLEA residential scenario without home-grown produce




Table 1: Exposure assessment parameters for residential scenario without home-grown produce – inputs for CLEA model

Parameter	Value	Justification
Land use	Residential without home-grown produce	Chosen land use
Receptor	Female child	Key generic assumption given in Box 3.1, SR3 ⁽⁵⁾
Building	Small terraced house	Key generic assumption given in Box 3.1, SR3 ⁽⁵⁾ . Small, two-storey terraced house chosen, as it is the most conservative residential building type in terms of protection from vapor intrusion (Section 3.4.6, SR3) ⁽⁵⁾
Soil type	Sandy loam	Most common UK soil type (Section 4.3.1, from Table 3.1, SR3) ⁽⁵⁾
Start age class (AC)	1	Range of age classes corresponding to key generic assumption that the critical receptor is a young female child aged 0–6. From Box 3.1, SR3 ⁽⁵⁾
End AC	6	
SOM (%)	6	Representative of sandy loamy soil according to EA guidance note dated January 2009 entitled 'Changes We Have Made to the CLEA Framework Documents' ⁽¹³⁾
	1	To provide SAC for sites where SOM <6% as often observed by RSK
	2.5	
pH	7	Model default

Table 2: Residential without home-grown produce – modified receptor data

Parameter	Unit	Age class					
		1	2	3	4	5	6
Soil to skin adherence factor – (outdoor)	mg soil/cm ² skin	0.1	0.1	0.1	0.1	0.1	0.1
Justification		Table 3.5, SP1010 ⁽³⁾					
Inhalation rate	m ³ day ⁻¹	5.4	8.0	8.9	10.1	10.1	10.1
Justification		Mean value USEPA, 2011 ⁽¹²⁾ ; Table 3.2, SP1010 ⁽³⁾					
Notes: For cadmium , the exposure assessment for a residential land use is based on estimates representative of lifetime exposure AC1-18. This is because the TDI _{ora} and TDI _{inh} are based on considerations of the kidney burden accumulated over 50 years. It is therefore reasonable to consider exposure not just in childhood but averaged over a longer period. See the Environment Agency Science Report SC05002/ TOX 3 ⁽¹⁾ , Science Report SC050021/Cadmium SGV ⁽¹⁾ and the project report SP1010 ⁽³⁾ for more information.							

References

1. Environment Agency (2009), 'Science Reports SC050021 - SGV and TOX reports for: benzene, toluene, ethylbenzene, xylene, mercury, selenium, nickel, arsenic, cadmium, phenol, dioxins, furans and dioxin-like PCBs'; 'Supplementary information for the derivation of SGV for: benzene, toluene, ethylbenzene, xylene, mercury, selenium, nickel, arsenic, cadmium, phenol, dioxins, furans and dioxin-like PCBs', and 'Contaminants in soil: updated collation of toxicological data and intake values for humans: benzene, toluene, ethylbenzene, xylene, mercury, selenium, nickel, arsenic, cadmium, phenol, dioxins, furans and dioxin-like PCBs'. Available at: <https://www.gov.uk/government/publications/contaminants-in-soil-updated-collation-of-toxicological-data-and-intake-values-for-humans> and <https://www.gov.uk/government/publications/land-contamination-soil-guideline-values-sgvs> (accessed 4 February 2015)
2. Nathanial, C. P., McCaffrey, C., Ashmore, M., Cheng, Y., Gillet, A. G., Ogden, R. C. and Scott, D. (2009), *LQM/CIEH Generic Assessment Criteria for Human Health Risk Assessment*, second edition (Nottingham: Land Quality Press).
3. Contaminated Land: Applications in Real Environment (CL:AIRE) (2014). 'Development of Category 4 Screening Levels for Assessment of Land Affected by Contamination', Revision 2, DEFRA research project SP1010.
4. Department for Environment, Food and Rural Affairs (Defra) (2014), 'SP1010: Development of Category 4 Screening Levels for assessment of land affected by contamination – Policy Companion Document', Revision 2.
5. Environment Agency (2009), *Science Report – SC050021/SR3. Updated technical background to the CLEA model* (Bristol: Environment Agency).
6. Contaminated Land: Applications in Real Environment (CL:AIRE) (2014). 'Appendices C to H). DEFRA research project SP1010'.
7. Nathanial, C. P., McCaffrey, C., Gillet, A. G., Ogden, R. C. and Nathanial, J. F. (2015), *The LQM/CIEH S4ULs for Human Health Risk Assessment* (Nottingham: Land Quality Press).
8. Environment Agency (2009), *Human health toxicological assessment of contaminants in soil. Science Report – Final SC050021/SR2* (Bristol: Environment Agency).
9. Environment Agency (2009), *Science Report – SC050021/SR4 CLEA Software (version 1.05) Handbook* (Bristol: Environment Agency).
10. Environment Agency (2008), *Science Report SC050021/SR7. Compilation of Data for Priority Organic Pollutants for Derivation of Soil Guideline Values* (Bristol: Environment Agency).
11. CL:AIRE (2010), *Soil Generic Assessment Criteria for Human Health Risk Assessment* (London: CL:AIRE).
12. USEPA (2011), *Exposure factors handbook*, EPA/600/R-090/052F (Washington, DC: Office of Research and Development).
13. Environment Agency (2009), 'Changes made to the CLEA framework documents after the three-month evaluation period in 2008', released January 2009.
14. USEPA (2010). Hydrogen cyanide and cyanide salts. Integrated Risk Information Systems (IRIS) Chemical Assessment Summary. September 2010. <https://www.epa.gov/iris> (accessed 9 December 2015)

GENERIC ASSESSMENT CRITERIA FOR HUMAN HEALTH - RESIDENTIAL WITHOUT HOME-GROWN PRODUCE



Table 3
Human Health Generic Assessment Criteria by Pathway for Residential Scenario Without Home-Grown Produce

Compound	Notes	SAC Appropriate to Pathway SOM 1% (mg/kg)			Soil Saturation Limit (mg/kg)	SAC Appropriate to Pathway SOM 2.5% (mg/kg)			Soil Saturation Limit (mg/kg)	SAC Appropriate to Pathway SOM 6% (mg/kg)			Soil Saturation Limit (mg/kg)
		Oral	Inhalation	Combined		Oral	Inhalation	Combined		Oral	Inhalation	Combined	
Metals													
Arsenic	(a,b)	3.99E+01	5.26E+02	NR	NR	3.99E+01	5.26E+02	NR	NR	3.99E+01	5.26E+02	NR	NR
Barium	(b)	1.35E+03	NR	NR	NR	1.35E+03	NR	NR	NR	1.35E+03	NR	NR	NR
Beryllium		1.56E+02	1.72E+00	NR	NR	1.56E+02	1.72E+00	NR	NR	1.56E+02	1.72E+00	NR	NR
Boron		1.08E+04	5.20E+06	NR	NR	1.08E+04	5.20E+06	NR	NR	1.08E+04	5.20E+06	NR	NR
Cadmium	(a)	1.95E+02	4.88E+02	1.49E+02	NR	1.95E+02	4.88E+02	1.49E+02	NR	1.95E+02	4.88E+02	1.49E+02	NR
Chromium (III) - trivalent	(c)	1.98E+04	9.07E+02	NR	NR	1.98E+04	9.07E+02	NR	NR	1.98E+04	9.07E+02	NR	NR
Chromium (VI) - hexavalent	(a,d)	5.91E+01	2.06E+01	NR	NR	5.91E+01	2.06E+01	NR	NR	5.91E+01	2.06E+01	NR	NR
Copper		1.08E+04	1.41E+04	7.13E+03	NR	1.08E+04	1.41E+04	7.13E+03	NR	1.08E+04	1.41E+04	7.13E+03	NR
Lead	(a)	3.14E+02	NR	NR	NR	3.14E+02	NR	NR	NR	3.14E+02	NR	NR	NR
Elemental Mercury (Hg ⁰)	(d)	NR	2.41E-01	NR	4.31E+00	NR	5.74E-01	NR	1.07E+01	NR	1.25E+00	NR	2.58E+01
Inorganic Mercury (Hg ²⁺)		5.71E+01	3.63E+03	5.62E+01	NR	5.71E+01	3.63E+03	5.62E+01	NR	5.71E+01	3.63E+03	5.62E+01	NR
Methyl Mercury (Hg ⁴⁺)		1.80E+01	1.87E+01	9.16E+00	7.33E+01	1.80E+01	3.62E+01	1.20E+01	1.42E+02	1.80E+01	7.68E+01	1.46E+01	3.04E+02
Nickel	(d)	1.88E+02	1.81E+02	NR	NR	1.88E+02	1.81E+02	NR	NR	1.88E+02	1.81E+02	NR	NR
Selenium	(b)	4.31E+02	NR	NR	NR	4.31E+02	NR	NR	NR	4.31E+02	NR	NR	NR
Vanadium		1.17E+03	1.46E+03	NR	NR	1.17E+03	1.46E+03	NR	NR	1.17E+03	1.46E+03	NR	NR
Zinc	(b)	4.05E+04	3.63E+07	NR	NR	4.05E+04	3.63E+07	NR	NR	4.05E+04	3.63E+07	NR	NR
Cyanide (free)		4.03E+01	1.37E+04	4.02E+01	NR	4.03E+01	1.37E+04	4.02E+01	NR	4.03E+01	1.37E+04	4.02E+01	NR
Volatile Organic Compounds													
Benzene	(a)	7.36E+01	9.01E-01	8.90E-01	1.22E+03	7.36E+01	1.68E+00	1.64E+00	2.26E+03	7.36E+01	3.48E+00	3.33E+00	4.71E+03
Toluene		2.87E+04	9.08E+02	8.80E+02	8.69E+02	2.87E+04	2.00E+03	1.87E+03	1.92E+03	2.87E+04	4.55E+03	3.93E+03	4.36E+03
Ethylbenzene		1.29E+04	8.34E+01	8.29E+01	5.18E+02	1.29E+04	1.96E+02	1.93E+02	1.22E+03	1.29E+04	4.58E+02	4.42E+02	2.84E+03
Xylene - m		2.32E+04	8.25E+01	8.22E+01	6.25E+02	2.32E+04	1.95E+02	1.93E+02	1.47E+03	2.32E+04	4.56E+02	4.47E+02	3.46E+03
Xylene - o		2.32E+04	8.87E+01	8.83E+01	4.78E+02	2.32E+04	2.08E+02	2.06E+02	1.12E+03	2.32E+04	4.86E+02	4.76E+02	2.62E+03
Xylene - p		2.32E+04	7.93E+01	7.90E+01	5.76E+02	2.32E+04	1.86E+02	1.85E+02	1.35E+03	2.32E+04	4.36E+02	4.28E+02	3.17E+03
Total xylene		2.32E+04	7.93E+01	7.90E+01	6.25E+02	2.32E+04	1.86E+02	1.85E+02	1.47E+03	2.32E+04	4.36E+02	4.28E+02	3.46E+03
Methyl tertiary-Butyl ether (MTBE)		3.87E+04	1.04E+02	1.04E+02	2.04E+04	3.87E+04	1.69E+02	1.69E+02	3.31E+04	3.87E+04	3.21E+02	3.19E+02	6.27E+04
Trichloroethene		6.45E+01	1.72E-02	1.72E-02	1.54E+03	6.45E+01	3.59E-02	3.59E-02	3.22E+03	6.45E+01	7.98E-02	7.97E-02	7.14E+03
Tetrachloroethene		7.13E+02	1.79E-01	1.79E-01	4.24E+02	7.13E+02	4.02E-01	4.02E-01	9.51E+02	7.13E+02	9.21E-01	9.20E-01	2.18E+03
1,1,1-Trichloroethane		7.74E+04	9.01E+00	9.01E+00	1.43E+03	7.74E+04	1.84E+01	1.84E+01	2.92E+03	7.74E+04	4.04E+01	4.04E+01	6.39E+03
1,1,1,2-Tetrachloroethane		7.34E+02	1.54E+00	1.53E+00	2.60E+03	7.34E+02	3.56E+00	3.55E+00	6.02E+03	7.34E+02	8.29E+00	8.20E+00	1.40E+04
1,1,2,2-Tetrachloroethane		7.34E+02	3.92E+00	3.90E+00	2.67E+03	7.34E+02	8.04E+00	7.95E+00	5.46E+03	7.34E+02	1.76E+01	1.72E+01	1.20E+04
Carbon Tetrachloride		5.15E+02	2.58E-02	2.58E-02	1.52E+03	5.15E+02	5.65E-02	5.64E-02	3.32E+03	5.15E+02	1.28E-01	1.28E-01	7.54E+03
1,2-Dichloroethane		1.55E+01	9.20E-03	9.20E-03	3.41E+03	1.55E+01	1.33E-02	1.33E-02	4.91E+03	1.55E+01	2.28E-02	2.27E-02	8.43E+03
Vinyl Chloride		1.81E+00	7.73E-04	7.73E-04	1.36E+03	1.81E+00	1.00E-03	9.99E-04	1.76E+03	1.81E+00	1.53E-03	1.53E-03	2.69E+03
1,2,4-Trimethylbenzene		NR	5.58E+00	NR	4.74E+02	NR	1.29E+01	NR	1.16E+03	NR	2.69E+01	NR	2.76E+03
1,3,5-Trimethylbenzene	(e)	NR	NR	NR	2.30E+02	NR	NR	NR	5.52E+02	NR	NR	NR	1.30E+03
Semi-Volatile Organic Compounds													
Acenaphthene		7.64E+03	4.86E+04	6.60E+03	5.70E+01	7.64E+03	1.18E+05	7.17E+03	1.41E+02	7.64E+03	2.68E+05	7.43E+03	3.36E+02
Acenaphthylene		7.65E+03	4.59E+04	6.55E+03	8.61E+01	7.65E+03	1.11E+05	7.15E+03	2.12E+02	7.65E+03	2.53E+05	7.42E+03	5.06E+02
Anthracene		3.82E+04	1.53E+05	3.06E+04	1.17E+00	3.82E+04	3.77E+05	3.47E+04	2.91E+00	3.82E+04	8.76E+05	3.66E+04	6.96E+00
Benzo(a)anthracene		1.98E+01	2.47E+01	1.10E+01	1.71E+00	1.98E+01	4.37E+01	1.36E+01	4.28E+00	1.98E+01	6.26E+01	1.50E+01	1.03E+01
Benzo(a)pyrene	(a)	5.34E+00	3.51E+01	NR	9.11E-01	5.34E+00	3.77E+01	NR	2.28E+00	5.34E+00	3.89E+01	NR	5.46E+00
Benzo(b)fluoranthene		4.97E+00	1.93E+01	3.95E+00	1.22E+00	4.97E+00	2.13E+01	4.03E+00	3.04E+00	4.97E+00	2.22E+01	4.06E+00	7.29E+00
Benzo(g,h,i)perylene		4.38E+02	1.87E+03	3.55E+02	1.54E-02	4.38E+02	1.94E+03	3.58E+02	3.85E-02	4.38E+02	1.97E+03	3.59E+02	9.23E-02
Benzo(k)fluoranthene		1.31E+02	5.41E+02	1.06E+02	6.87E-01	1.31E+02	5.76E+02	1.07E+02	1.72E+00	1.31E+02	5.91E+02	1.07E+02	4.12E+00
Chrysene		3.95E+01	1.19E+02	2.97E+01	4.40E-01	3.95E+01	1.49E+02	3.12E+01	1.10E+00	3.95E+01	1.66E+02	3.19E+01	2.64E+00
Dibenzo(a,h)anthracene		3.95E-01	1.45E+00	3.10E-01	3.93E-03	3.95E-01	1.64E+00	3.18E-01	9.82E-03	3.95E-01	1.74E+00	3.22E-01	2.36E-02
Fluoranthene		1.59E+03	3.83E+04	1.53E+03	1.89E+01	1.59E+03	8.87E+04	1.56E+03	4.73E+01	1.59E+03	1.83E+05	1.58E+03	1.13E+02

GENERIC ASSESSMENT CRITERIA FOR HUMAN HEALTH - RESIDENTIAL WITHOUT HOME-GROWN PRODUCE



Table 3

Human Health Generic Assessment Criteria by Pathway for Residential Scenario Without Home-Grown Produce

Compound	Notes	SAC Appropriate to Pathway SOM 1% (mg/kg)			Soil Saturation Limit (mg/kg)	SAC Appropriate to Pathway SOM 2.5% (mg/kg)			Soil Saturation Limit (mg/kg)	SAC Appropriate to Pathway SOM 6% (mg/kg)			Soil Saturation Limit (mg/kg)
		Oral	Inhalation	Combined		Oral	Inhalation	Combined		Oral	Inhalation	Combined	
Fluorene		5.09E+03	6.20E+03	2.80E+03	3.09E+01	5.09E+03	1.53E+04	3.82E+03	7.65E+01	5.09E+03	3.62E+04	4.47E+03	1.83E+02
Indeno(1,2,3-cd)pyrene		5.65E+01	2.12E+02	4.46E+01	6.13E-02	5.65E+01	2.38E+02	4.56E+01	1.53E-01	5.65E+01	2.50E+02	4.60E+01	3.68E-01
Naphthalene		2.50E+03	2.33E+01	2.31E+01	7.64E+01	2.50E+03	5.58E+01	5.46E+01	1.83E+02	2.50E+03	1.31E+02	1.25E+02	4.32E+02
Phenanthrene		1.58E+03	7.17E+03	1.30E+03	3.60E+01	1.58E+03	1.76E+04	1.45E+03	8.96E+01	1.58E+03	4.07E+04	1.52E+03	2.14E+02
Pyrene		3.82E+03	8.79E+04	3.66E+03	2.20E+00	3.82E+03	2.04E+05	3.75E+03	5.49E+00	3.82E+03	4.23E+05	3.79E+03	1.32E+01
Phenol		6.48E+04	4.58E+02	4.55E+02	2.42E+04	6.48E+04	6.95E+02	6.88E+02	3.81E+04	6.48E+04	1.19E+03	1.17E+03	7.03E+04
Total Petroleum Hydrocarbons													
Aliphatic hydrocarbons EC ₅ -EC ₆		3.23E+05	4.24E+01	4.24E+01	3.04E+02	3.23E+05	7.79E+01	7.79E+01	5.58E+02	3.23E+05	1.61E+02	1.61E+02	1.15E+03
Aliphatic hydrocarbons >EC ₆ -EC ₈		3.23E+05	1.04E+02	1.04E+02	1.44E+02	3.23E+05	2.31E+02	2.31E+02	3.22E+02	3.23E+05	5.29E+02	5.29E+02	7.36E+02
Aliphatic hydrocarbons >EC ₈ -EC ₁₀		6.45E+03	2.68E+01	2.68E+01	7.77E+01	6.45E+03	6.55E+01	6.53E+01	1.90E+02	6.45E+03	1.56E+02	1.55E+02	4.51E+02
Aliphatic hydrocarbons >EC ₁₀ -EC ₁₂		6.45E+03	1.33E+02	1.32E+02	4.75E+01	6.45E+03	3.31E+02	3.27E+02	1.18E+02	6.45E+03	7.93E+02	7.67E+02	2.83E+02
Aliphatic hydrocarbons >EC ₁₂ -EC ₁₆		6.45E+03	1.11E+03	1.06E+03	2.37E+01	6.45E+03	2.78E+03	2.42E+03	5.91E+01	6.45E+03	6.67E+03	4.37E+03	1.42E+02
Aliphatic hydrocarbons >EC ₁₆ -EC ₃₅	(b)	6.50E+04	NR	NR	8.48E+00	9.25E+04	NR	NR	2.12E+01	1.11E+05	NR	NR	5.09E+01
Aliphatic hydrocarbons >EC ₃₅ -EC ₄₄	(b)	6.50E+04	NR	NR	8.48E+00	9.25E+04	NR	NR	2.12E+01	1.11E+05	NR	NR	5.09E+01
Aromatic hydrocarbons >EC ₈ -EC ₁₀		2.58E+03	4.74E+01	4.72E+01	6.13E+02	2.58E+03	1.16E+02	1.15E+02	1.50E+03	2.58E+03	2.77E+02	2.69E+02	3.58E+03
Aromatic hydrocarbons >EC ₁₀ -EC ₁₂		2.58E+03	2.58E+02	2.52E+02	3.64E+02	2.58E+03	6.39E+02	5.94E+02	8.99E+02	2.58E+03	1.52E+03	1.24E+03	2.15E+03
Aromatic hydrocarbons >EC ₁₂ -EC ₁₆		2.58E+03	2.85E+03	1.80E+03	1.69E+02	2.58E+03	7.07E+03	2.30E+03	4.19E+02	2.58E+03	1.68E+04	2.48E+03	1.00E+03
Aromatic hydrocarbons >EC ₁₆ -EC ₂₁	(b)	1.86E+03	NR	NR	5.37E+01	1.90E+03	NR	NR	1.34E+02	1.92E+03	NR	NR	3.21E+02
Aromatic hydrocarbons >EC ₂₁ -EC ₃₅	(b)	1.93E+03	NR	NR	4.83E+00	1.93E+03	NR	NR	1.21E+01	1.93E+03	NR	NR	2.90E+01
Aromatic hydrocarbons >EC ₃₅ -EC ₄₄	(b)	1.93E+03	NR	NR	4.83E+00	1.93E+03	NR	NR	1.21E+01	1.93E+03	NR	NR	2.90E+01

Notes:

EC - equivalent carbon. GrAC - groundwater assessment criteria. SAC - soil assessment criteria.

The CLEA model output is colour coded depending upon whether the soil saturation limit has been exceeded.

	Calculated SAC exceeds soil saturation limit and may significantly affect the interpretation of any exceedances as the contribution of the indoor and outdoor vapour pathway to total exposure is >10%.
	Calculated SAC exceeds soil saturation limit but the exceedance will not affect the SAC significantly as the contribution of the indoor and outdoor vapour pathway to total exposure is <10%.
	Calculated SAC does not exceed the soil saturation limit.

The SAC for organic compounds are dependant upon soil organic matter (SOM) (%) content. To obtain SOM from total organic carbon (TOC) (%) divide by 0.58. 1% SOM is 0.58% TOC. DL Rowell Soil Science: Methods and Applications, Longmans, 1994.

SAC for TPH fractions, PAHs naphthalene, acenaphthene and acenaphthylene, BTEX and trimethylbenzene compounds were produced using an attenuation factor for the indoor air inhalation pathway of 10 to reduce conservatism associated with the vapour inhalation pathway (Section 10.1.1, SR3)

(a) SAC for arsenic, benzene, benzo(a)pyrene, cadmium, chromium VI and lead are derived using the C4SL toxicology data.

(b) SAC for boron and selenium should not include the inhalation pathway as no expert group HCV has been derived; aliphatic and aromatic hydrocarbons >EC16 should not include inhalation pathway due to their non-volatile nature and inhalation exposure being minimal (oral, dermal and inhalation exposure is compared to the oral HCV); arsenic should only be based on oral contribution (rather than combined) owing to the relative small contribution from inhalation in accordance with the SGV report. The Oral SAC should be adopted for zinc and benzo(a)pyrene.

(c) SAC for CrIII should be based on the lower of the oral and inhalation SAC (see LQM/CIEH 2015 Section 6.8)

(d) SAC for elemental mercury, chromium VI and nickel should be based on the inhalation pathway only.

(e) SAC for 1,3,5-trimethylbenzene is not recorded owing to the lack of toxicological data, SAC for 1,2,4 trimethylbenzene may be used.

GENERIC ASSESSMENT CRITERIA FOR HUMAN HEALTH - RESIDENTIAL WITHOUT HOME-GROWN PRODUCE			
<div>Table 4</div> <div>Human health generic assessment criteria for residential without home-grown produce</div>			
<div>RSK</div>			
Compound	SAC for Soil SOM 1% (mg/kg)	SAC for Soil SOM 2.5% (mg/kg)	SAC for Soil SOM 6% (mg/kg)
Metals			
Arsenic	40	40	40
Barium	1,300	1,300	1,300
Beryllium	1.7	1.7	1.7
Boron	11,000	11,000	11,000
Cadmium	149	149	149
Chromium (III) - trivalent	910	910	910
Chromium (VI) - hexavalent	21	21	21
Copper	7,100	7,100	7,100
Lead	310	310	310
Elemental Mercury (Hg ⁰)	0.2	0.6	1.2
Inorganic Mercury (Hg ²⁺)	56	56	56
Methyl Mercury (Hg ⁴⁺)	9	12	15
Nickel	180	180	180
Selenium	430	430	430
Vanadium	1,200	1,200	1,200
Zinc	40,000	40,000	40,000
Cyanide (free)	40	40	40
Volatile Organic Compounds			
Benzene	0.9	1.6	3.3
Toluene	900 (869)	1,900	3,900
Ethylbenzene	80	190	440
Xylene - m	80	190	450
Xylene - o	90	210	480
Xylene - p	80	180	430
Total xylene	80	180	430
Methyl tertiary-Butyl ether (MTBE)	100	170	320
Trichloroethene	0.02	0.04	0.08
Tetrachloroethene	0.2	0.4	0.9
1,1,1-Trichloroethane	9.0	18.4	40.4
1,1,1,2-Tetrachloroethane	1.5	3.5	8.2
1,1,2,2-Tetrachloroethane	3.9	8.0	17.2
Carbon Tetrachloride	0.026	0.056	0.128
1,2-Dichloroethane	0.009	0.013	0.023
Vinyl Chloride	0.0008	0.0010	0.0015
1,2,4-Trimethylbenzene	5.6	12.9	26.9
1,3,5-Trimethylbenzene	NR	NR	NR
Semi-Volatile Organic Compounds			
Acenaphthene	6,600 (57)	7,200	7,400
Acenaphthylene	6,600 (86)	7,200	7,400
Anthracene	31,000 (1.17)	35,000	37,000
Benzo(a)anthracene	11.0	13.6	15.0
Benzo(a)pyrene	5.3	5.3	5.3
Benzo(b)fluoranthene	4.0	4.0	4.1
Benzo(g,h,i)perylene	355	358	359
Benzo(k)fluoranthene	106	107	107
Chrysene	30	31	32
Dibenzo(a,h)anthracene	0.31	0.32	0.32
Fluoranthene	1,500	1,600	1,600
Fluorene	2,800 (31)	3,800 (77)	4,500 (183)
Indeno(1,2,3-cd)pyrene	45	46	46
Naphthalene	23	55	125
Phenanthrene	1,300 (36)	1,450	1,520
Pyrene	3,700	3,800	3,800
Phenol	440*	688	1,170
Total Petroleum Hydrocarbons			
Aliphatic hydrocarbons EC ₅ -EC ₆	42	78	161
Aliphatic hydrocarbons >EC ₆ -EC ₈	100	230	530
Aliphatic hydrocarbons >EC ₈ -EC ₁₀	27	65	155
Aliphatic hydrocarbons >EC ₁₀ -EC ₁₂	130 (48)	330 (118)	770 (283)
Aliphatic hydrocarbons >EC ₁₂ -EC ₁₆	1,100 (24)	2,400 (59)	4,400 (142)
Aliphatic hydrocarbons >EC ₁₆ -EC ₃₅	65,000 (8)	92,000 (21)	111,000
Aliphatic hydrocarbons >EC ₃₅ -EC ₄₄	65,000 (8)	92,000 (21)	111,000
Aromatic hydrocarbons >EC ₈ -EC ₁₀	47	115	269
Aromatic hydrocarbons >EC ₁₀ -EC ₁₂	300	600	1,200
Aromatic hydrocarbons >EC ₁₂ -EC ₁₆	1,800 (169)	2,300 (419)	2,500
Aromatic hydrocarbons >EC ₁₆ -EC ₂₁	1,900	1,900	1,900
Aromatic hydrocarbons >EC ₂₁ -EC ₃₅	1,900	1,900	1,900
Aromatic hydrocarbons >EC ₃₅ -EC ₄₄	1,900	1,900	1,900
Minerals			
Asbestos	Stage 1 test – No asbestos detected with ID; Stage 2 test - <0.001% dry weight (exceedance of either equates to an exceedance of the GAC) ¹		
Notes:			
* - Generic assessment criteria not calculated owing to low volatility of substance and therefore no pathway, or an absence of toxicological data.			
NR - SAC for 1,3,5-trimethylbenzene is not recorded owing to the lack of toxicological data, SAC for 1,2,4 trimethylbenzene may be used			
EC - equivalent carbon. SAC - soil assessment criteria.			
¹ LOD for weight of asbestos per unit weight of soil calculated on a dry weight basis using PLM, handpicking and gravimetry.			
The SAC for organic compounds are dependent on Soil Organic Matter (SOM) (%) content. To obtain SOM from total organic carbon (TOC) (%) divide by 0.58.			
1% SOM is 0.58% TOC. DL Rowell Soil Science: Methods and Applications, Longmans, 1994.			
SAC for TPH fractions, PAHs naphthalene, acenaphthene and acenaphthylene, BTEX and trimethylbenzene compounds were produced using an attenuation factor for the indoor air inhalation pathway of 10 to reduce conservatism associated with the vapour inhalation pathway, section 10.1.1, SR3.			
(VALUE IN BRACKETS)			
RSK has adopted an approach for petroleum hydrocarbons in accordance with LQM/CIEH whereby the concentration modelled for each petroleum hydrocarbon fraction has been tabulated as the SAC with the corresponding solubility or vapour saturation limits given in brackets.			



APPENDIX K

RSK GROUNDWATER GACS

GENERIC ASSESSMENT CRITERIA FOR CONTROLLED WATERS

Protection of the water environment

The water environment in the United Kingdom is protected under a number of regulatory regimes. The relevant environmental regulator is consulted where there may be a risk that pollution of 'controlled waters' may occur or may have occurred in the past.

The term 'controlled waters' refers to coastal waters, inland freshwaters and groundwater. The EU Water Framework Directive (WFD) (2000/60/EC) is implemented via domestic regulations and guidance, covering aspects of groundwater and surface water protection as well as drinking water supply policy. Domestic legislation and guidance will vary across the United Kingdom. Therefore, the relevant legislation for England, Wales, Northern Ireland and Scotland should be reviewed, alongside guidance provided by the Environment Agency (EA), Natural Resource Wales (NRW), the Scottish Environmental Protection Agency (SEPA) or the Northern Ireland Environment Agency (NIEA), as appropriate.

The main objectives of the protection and remediation of groundwater under threat from land contamination are set out within "The Environment Agency's approach to groundwater protection", version 1.0 (March 2017)⁽¹⁾ and the associated guidance "Land contamination groundwater compliance points: quantitative risk assessments (March 2017)^(1a) that have replaced the previous guidance document "Groundwater Principles and Practice (GP3)". When assessing risks to groundwater, the following need to be considered:

- Where pollutants have not yet entered groundwater, all necessary and reasonable measures must be taken to:
 - **prevent** the input of **hazardous** substances into groundwater (see description of hazardous substances below)
 - **limit** the entry of other (non-hazardous) pollutants into groundwater to avoid pollution, deterioration in the status of groundwater bodies and to prevent sustained, upward trends in pollutant concentrations in groundwater.
- Where pollutants have already entered groundwater, the priority is to take all necessary and reasonable measures to:
 - **minimise** further entry of "contaminants" where there is a defined source
 - **limit the pollution** of groundwater or any effect on the status of the groundwater body from the future expansion of the 'plume', if necessary, by actively reducing its extent.

Within the context of groundwater risk assessments on sites affected by land contamination, "reasonable" means feasible without involving disproportionate costs. What costs are "disproportionate" depends on site-specific circumstances, which may include:

- Considerations of technical feasibility such as identified by the remedial options appraisal, this may be due to the distribution or nature of the contamination and the available remedial methods to treat the identified contamination;
- Sustainability considerations.

DEFINITIONS AND SUBSTANCE CLASSIFICATIONS

Risks to surface waters:

When assessing risks to surface waters, the following list of definitions should be understood:

Priority substances (PS) are harmful substances originally identified under the Water Framework Directive (WFD) 2000/60/EC as substances ‘presenting a significant risk to or via the aquatic environment’ at a European level. Member States are required to incorporate the identified **PS** into their country-wide monitoring programmes. There are currently 33 **PS** defined within the Priority Substances Directive (2013/39/EU; Annex 1), with a further 12 additional substances due to come into force from 22 December 2018. Directive 2013/39/EU has been transposed into domestic legislation for England and Wales by The Water Framework Directive (Standards and Classification) Directions (England and Wales) 2015.

Under the umbrella of **PS**, there is a sub-set of substances identified as being “hazardous”, and these are referred to as **Priority hazardous substances (PHS)**. The list of **PHS** is defined at EU level within the Priority Substances Directive (2013/39/EU). The WFD defines hazardous substances as ‘substances (or groups of substances) that are toxic, persistent and liable to bio-accumulate, and other substances or groups of substances that give rise to an equivalent level of concern.’ There are currently 15 **PHS**, with a further 6 additional substances due to come into force from 22 December 2018.

There is also another group of substances defined at EU level and which are referred to as **other pollutants (OP)** in Directive 2013/39/EU. These are additional substances which although not **priority substances**, have EQS which are identical to those laid down in the legislation which applied prior to 13 January 2009 (Directive 2008/105/EU). The **OP** are listed along with the **priority substance (PS)** within the Priority Substances Directive (2013/39/EU), and their associated EQS are also listed therein. There are 6 **OP** defined within the Priority Substances Directive (2013/39/EU).

In addition to the EU level substances, there are also a group of pollutants defined at a Member State level, referred to as **Specific pollutants (SP)**. These substances are pollutants which are released in significant quantities into water bodies in each of the individual European Member States. Under the WFD, Member States are required to set their own EQS for these substances. An indicative list of **SP** is given in Annex VIII of the WFD. Many of the substances categorised as **SP** in the UK were formerly List 2 substances under the old Groundwater Directive (80/68/EEC). The **SP** are defined within Part 2 (Table 1) of The Water Framework Directive (Standards and Classification) Directions (England and Wales) 2015.

Risks to groundwater:

When assessing risks to groundwater, the following definitions should be understood:

Under the requirements of the Groundwater Daughter Directive (2006/118/EU), the UK has published a list of substances it considers to be **hazardous substances** with respect to groundwater. In their advisory capacity to the government, this list has been derived by the UK Joint Agencies Groundwater Directive Advisory Group (JAGDAG), of which the Environment Agency is a member. The JAGDAG list of **hazardous substances** was published in January 2017 and the Environment Agency will use the updated list of hazardous substances from this date for all new activities that may lead to the discharge of hazardous substances to groundwater. The list is extensive and can be found in full at:

<https://www.wfduk.org/stakeholders/jagdag>

Selecting the appropriate assessment criteria

When assessing the risks to controlled waters, various assessment criteria apply, depending on the nature of the assessment and the conceptual site model.

Where a surface water body is involved, then Environmental Quality Standards (EQS) are the relevant assessment criteria as they are designed to be protective of surface water ecology.

Where a public water supply or a Principal aquifer is involved, then the standards defined in The Water Supply (Water Quality) Regulations⁽²⁾ are the primary source of assessment criteria. The Private Water Supplies Regulations⁽³⁾ may also be applicable in some cases. For instances where there are no UK assessment criteria, then the World Health Organisation (WHO) drinking water guidelines⁽⁴⁾ may be used.

This appendix presents the generic assessment criteria (GAC) that RSK considers suitable for assessing risks to controlled waters for our most commonly encountered determinants. A full list of EQS for England and Wales are included in The Water Framework Directive (Standards and Classification) Directions (England and Wales) 2015.

The RSK GAC for controlled waters are presented in **Table 1** and **Table 2**. In line with the Environment Agency's Remedial Targets Methodology, the GAC for controlled waters are termed 'target concentrations'.

The appropriate target concentrations should be selected with consideration to:

- the site conceptual model (i.e. the receptor at potential risk);
- whether the substance is already present in groundwater at the site;
- whether or not the substance is classified as a priority hazardous substance under the Priority Substances Directive (2013/39/EC) (see above), or as a hazardous substance according to the current list of JAGDAG determinations⁽⁵⁾; and
- background concentrations in the aquifer (if applicable).

It is important to remember that the WFD and Environment Agency guidance^(1 & 1a) support a sustainable, risk-based approach be applied to groundwater contamination. Exceedance of any target concentration does not necessarily imply that an unacceptable risk exists or that remediation is inevitably required.

Target concentrations shaded in green are <u>statutory values</u>	Target concentrations shaded in orange are <u>non-statutory values</u>
---	--

Note: Units µg/l throughout (unless otherwise stated)

Table 1: Target concentrations for controlled waters (excluding TPH CWG fractions)

Substance classification		Determinant	Target concentrations (µg/l)			
Groundwater receptors ⁽⁵⁾	Surface water receptors ⁽⁶⁾		Minimum reporting value	UK drinking water standard (or best equivalent)	EQS or best equivalent	
					Freshwater	Transitional (estuaries) and coastal waters
Metals & other inorganics						
Hazardous substance	Specific pollutant	Arsenic	-	10 ⁽²⁾	50 ^(6a)	25 ^(6a)
Non-hazardous pollutant	Priority substance	Cadmium	0.1 ⁽⁷⁾	5 ⁽²⁾	≤0.08, 0.08, 0.09, 0.15, 0.25 ^(6b)	0.2 ^(6a)
(Not determined)	-	Chromium (total)	-	50 ⁽²⁾	8.1 Sum values for chromium III and VI	-
(None	Specific pollutant	Chromium (III)	-	Use value for total chromium	4.7 ^(6a)	-
Hazardous substance	Specific pollutant	Chromium (VI)			3.4 ^(6a)	0.6 ^(6a)

Substance classification		Determinant	Target concentrations (µg/l)			
Groundwater receptors ⁽⁵⁾	Surface water receptors ⁽⁶⁾		Minimum reporting value	UK drinking water standard (or best equivalent)	EQS or best equivalent	
					Freshwater	Transitional (estuaries) and coastal waters
(Not determined)	Specific pollutant	Copper	-	2,000 ⁽²⁾	1 bioavailable ^(6a)	3.76 dissolved, where DOC ≤1mg/l ^(6a)
						3.76µg/l + (2.677µg/l x ((DOC/2) – 0.5µg/l)) dissolved, where DOC >1mg/l ^(6a)
Hazardous substance	Priority substance	Lead	-	10 ⁽²⁾	1.2 bioavailable ^(6a)	1.3 ^(6a)
Hazardous substance	Priority hazardous substance	Mercury	0.01 ⁽⁷⁾	1 ⁽²⁾	0.07 ^(6c)	0.07 ^(6c)
Non-hazardous pollutant	Priority substance	Nickel	-	20 ⁽²⁾	4.0 bioavailable ^(6a)	8.6 ^(6a)
Non-hazardous pollutant	-	Selenium	-	10 ⁽²⁾	-	-
Non-hazardous pollutant	Specific pollutant	Zinc	-	3,000 ⁽⁸⁾	10.9 bioavailable ^(6a)	6.8 dissolved ^(6a)
None	Specific pollutant	Iron	-	200 ⁽²⁾	1000 ^(6a) *1	1000 ^(6a))*1
None	Specific pollutant	Manganese	-	50 ⁽²⁾ (0.05mg/l)	123 bioavailable ^(6a) (0.123mg/l)	-
(Not determined)	-	Aluminium	-	200 ⁽²⁾	-	-

Substance classification		Determinant	Target concentrations (µg/l)			
Groundwater receptors ⁽⁵⁾	Surface water receptors ⁽⁶⁾		Minimum reporting value	UK drinking water standard (or best equivalent)	EQS or best equivalent	
					Freshwater	Transitional (estuaries) and coastal waters
Hazardous substance	Priority hazardous substance	Tributyltin compounds (Tributyltin-cation)	0.001 ⁽⁷⁾	-	0.0002 ^(6a)	0.0002 ^(6a)
(Not determined)	-	Sodium	-	200,000 ⁽²⁾ (200 mg/l)	-	-
Non-hazardous pollutant	Specific pollutant	Cyanide (Hydrogen cyanide)	-	50 ⁽²⁾ (0.05 mg/l)	1 ^(6a) (0.001 mg/l)	1 ^(6a) (0.001 mg/l)
Non-hazardous pollutant	-	Total ammoniacal nitrogen ^s	-	500 ⁽²⁾ (0.5 mg/l) as NH ₄ (472 expressed as NH ₃ ; 389 expressed as N)	300 ^(6f) (0.3 mg/l) as N (364 expressed as NH ₃ ; 386 expressed as NH ₄)	-
Non-hazardous pollutant	Specific pollutant	Ammonia un-ionised (equilibrium ratio calculated) (NH ₃)	-	-	-	21 ^(6a) (0.021 mg/l)
Non-hazardous pollutant	Specific pollutant	Chlorine	-	-	2 ^(6a) (0.002 mg/l)	10 ^(6d) (0.01 mg/l)
(Not determined)	-	Chloride	-	250,000 ⁽²⁾ (250 mg/l)	-	-
(Not determined)	-	Sulphate	-	250,000 ⁽²⁾ (250 mg/l)	-	-
(Not determined)	-	Nitrate (as NO ₃)	-	50,000 ⁽²⁾ (50 mg/l)	-	-

Substance classification		Determinant	Target concentrations (µg/l)			
Groundwater receptors ⁽⁵⁾	Surface water receptors ⁽⁶⁾		Minimum reporting value	UK drinking water standard (or best equivalent)	EQS or best equivalent	
					Freshwater	Transitional (estuaries) and coastal waters
(Not determined)	-	Nitrite (as NO ₂)	-	500 ⁽²⁾ (0.5 mg/l)	10 ⁽⁹⁾ (0.01 mg/l)	-
Volatile organic compounds (VOC)						
Non-hazardous pollutant	Other pollutant	Tetrachloroethene (tetrachloroethylene; PCE)	0.1 ⁽⁷⁾	10 ⁽²⁾ sum of TCE and PCE	10 ^(6a)	10 ^(6a)
Hazardous substance	Other pollutant	Trichloroethene (trichloroethylene; TCE)	0.1 ⁽⁷⁾		10 ^(6a)	10 ^(6a)
None	Specific pollutant	Tetrachloroethane	-	-	140 ^(6a)	-
Hazardous substance	Other pollutant	Carbon tetrachloride (tetrachloromethane)	0.1 ⁽⁷⁾	3.0 ⁽²⁾	12 ^(6a)	12 ^(6a)
Non-hazardous pollutant	Priority substance	1,2-Dichloroethane	1.0 ⁽⁷⁾	3.0 ⁽²⁾	10 ^(6a)	10 ^(6a)
Non-hazardous pollutant	-	1.2-Dichloroethene (DCE) sum of cis and trans	-	50.0 ⁽⁴⁾	-	-
Hazardous substance	-	Vinyl chloride (chloroethene)	-	0.5 ⁽²⁾	-	-
Non-hazardous pollutant	Priority substance	Dichloromethane	-	20 ⁽⁴⁾	20 ^(6a)	20 ^(6a)
Non-hazardous pollutant	Priority substance	Trichlorobenzenes	0.01 ⁽⁷⁾	-	0.4 ^(6a)	0.4 ^{((6a)}

Substance classification		Determinant	Target concentrations (µg/l)			
Groundwater receptors ⁽⁵⁾	Surface water receptors ⁽⁶⁾		Minimum reporting value	UK drinking water standard (or best equivalent)	EQS or best equivalent	
					Freshwater	Transitional (estuaries) and coastal waters
Hazardous substance	Priority substance	Trichloromethane (Chloroform)	0.1 ⁽⁷⁾	100 ^(2a) (sum of trihalomethanes – chloroform, bromform, dibromochloromethane, bromodichloromethane)	2.5 ^(6a)	2.5 ^(6a)
(Not determined)	-	Bromoform	-		-	-
(Not determined)	-	Dibromochloromethane	-		-	-
(Not determined)	-	Bromodichloromethane	-		-	-
Non-hazardous pollutant	Priority hazardous substance	Di(2-ethylhexyl) phthalate (bis(2-ethylhexyl) phthalate, DEHP)	-	8 ⁽⁴⁾	1.3 ^(6a)	1.3 ^(6a)
None	Specific pollutant	Benzyl butyl phthalate	-	-	7.5 ^(6a)	0.75 ^(6e)
Hazardous substance	Priority hazardous substance	Hexachlorobutadiene (as a pesticide, but reported in a VOC suite)	0.005 ⁽⁷⁾	0.1 ⁽²⁾	0.6 ^(6c)	0.6 ^(6c)
Semi-volatile organic compounds (SVOC)						
(Not determined)	-	Acenaphthylene (Aro EC12-EC16)	-	-	5.8 ⁽¹⁰⁾	
Hazardous substance	Priority hazardous substance	Anthracene (Aro EC16-EC21)	-	-	0.1 ^(6a)	0.1 ^(6a)

Substance classification		Determinant	Target concentrations (µg/l)			
Groundwater receptors ⁽⁵⁾	Surface water receptors ⁽⁶⁾		Minimum reporting value	UK drinking water standard (or best equivalent)	EQS or best equivalent	
					Freshwater	Transitional (estuaries) and coastal waters
Non-hazardous pollutant	Priority substance	Naphthalene (Aro EC10-EC12)	-	-	2 ^(6a)	2 ^(6a)
Hazardous substance	Priority substance	Fluoranthene (Aro EC21-EC35) not used as an indicator for this EC band	-	-	0.0063 ^(6a)	0.0063 ^(6a)
Hazardous substance(s)	Priority hazardous substance(s)	Benzo(a)pyrene (Aro EC21-EC35)	-	0.01 ⁽²⁾	0.00017 ^(6a)	0.00017 ^(6a)
		Benzo(b)fluoranthene (Aro EC21-EC35)	-	0.1 ⁽²⁾ sum of the concentration of the four specified compounds	No EQS for these substances. B(a)P should be used as the indicator compound instead.	
		Benzo(k)fluoranthene (Aro EC21-EC35)	-			
		Benzo(g,h,i)perylene (Aro EC21-EC35)	-			
		Indeno(1,2,3-cd) pyrene (Aro EC21-EC35)	-			
Non-hazardous pollutant	Specific pollutant	Phenol	-	-	7.7 ^(6a)	7.7 ^(6a)
Hazardous substance	Specific pollutant	2,4-Dichlorophenol	0.1 ⁽⁷⁾	-	4.2 ^(6a)	0.42 ^(6a)

Substance classification		Determinant	Target concentrations (µg/l)			
Groundwater receptors ⁽⁵⁾	Surface water receptors ⁽⁶⁾		Minimum reporting value	UK drinking water standard (or best equivalent)	EQS or best equivalent	
					Freshwater	Transitional (estuaries) and coastal waters
Hazardous substance	Priority substance	Pentachloro-phenol (PCP) (as a pesticide, but reported in an SVOC suite)	0.1 ⁽⁷⁾	0.1 ⁽²⁾	0.4 ^(6a)	0.4 ^(6a)
Petroleum hydrocarbons						
Hazardous substance	-	Total petroleum hydrocarbons	-	See Table 2 for individual (non-statutory) TPH CWG fractions with respect to drinking water receptors	See individual risk driving compounds (i.e. BTEX and PAH) for specific EQS	
Hazardous substance	Priority substance	Benzene (Aro EC5-EC7)	1 ⁽⁷⁾	1 ⁽²⁾	10 ^(6a)	8 ^(6a)
Hazardous substance	Specific pollutant	Toluene (Aro EC7-EC8)	4 ⁽⁷⁾	700 ⁽⁴⁾	74 ^(6a)	74 ^(6a)
Hazardous substance	-	Ethylbenzene (Aro EC8-EC10)	-	300 ⁽⁴⁾	300 ⁽¹¹⁾	-
(Not determined)	-	Xylenes (Aro EC8-EC10)	3 ⁽⁷⁾	500 ⁽⁴⁾	30 ⁽¹¹⁾	-
Non-hazardous pollutant	-	Methyl tertiary butyl ether (MTBE)	-	15 ⁽¹²⁾	-	

Substance classification		Determinant	Target concentrations (µg/l)			
Groundwater receptors ⁽⁵⁾	Surface water receptors ⁽⁶⁾		Minimum reporting value	UK drinking water standard (or best equivalent)	EQS or best equivalent	
					Freshwater	Transitional (estuaries) and coastal waters
Pesticides, fungicides, insecticides and herbicides						
(Not determined) – assume to be Hazardous Substance	-	Total pesticides	-	0.5 ⁽²⁾	-	-
(Not determined) - assume to be Hazardous Substance	-	Other individual pesticides (unless otherwise detailed below)	-	0.1 ⁽²⁾	-	-
Hazardous substance(s)	Other pollutant (Cyclodiene pesticides)	Aldrin	0.003 ⁽⁷⁾	0.03 ⁽²⁾	0.01 ^(6a) (sum of all four)	0.005 ^(6a) (sum of all four)
		Dieldrin	0.003 ⁽⁷⁾	0.03 ⁽²⁾		
		Endrin	0.003 ⁽⁷⁾	0.1 ^(2b) ('other individual pesticide')		
		Isodrin* ²	0.003 ⁽⁷⁾	0.1 ^(2b) ('other individual pesticide')		
Hazardous substance	Other pollutant	DDT (total)	0.002 ⁽⁷⁾	0.1 ⁽²⁾ ('other individual pesticide')	0.025 ^(6a)	0.025 ^(6a)
Hazardous substance	Specific pollutant	Carbendazim	-	0.1 ⁽²⁾ ('other individual pesticide')	0.15 ^(6a)	-

Substance classification		Determinant	Target concentrations (µg/l)			
Groundwater receptors ⁽⁵⁾	Surface water receptors ⁽⁶⁾		Minimum reporting value	UK drinking water standard (or best equivalent)	EQS or best equivalent	
					Freshwater	Transitional (estuaries) and coastal waters
Hazardous substance	Specific pollutant	Chlorothalonil	-	0.1 ⁽²⁾ (‘other individual pesticide’)	0.035 ^(6a)	-
Hazardous substance	Specific pollutant (until 22/12/18, after which it becomes a Priority substance)	Cypermethrin	-	0.1 ⁽²⁾ (‘other individual pesticide’)	0.0001 ^(6a) From 22/12/18: 8.0E ^{-5(6a)}	0.0001 ^(6a) From 22/12/18: 8.0E ^{-6(6a)}
Hazardous substance	Specific pollutant	Dimethoate	0.01 ⁽⁷⁾	0.1 ⁽²⁾ (‘other individual pesticide’)	0.48 ^(6a)	0.48 ^(6a)
(Not determined)	Specific pollutant	Glyphosate	-	0.1 ⁽²⁾ (‘other individual pesticide’)	196 ^(6a)	196 ^(6a)
Hazardous substance	Specific pollutant	Linuron	-	0.1 ⁽²⁾ (‘other individual pesticide’)	0.5 ^(6a)	0.5 ^(6a)
Non-hazardous pollutant	Specific pollutant	Mecoprop	0.04 ⁽⁷⁾	0.1 ⁽²⁾ (‘other individual pesticide’)	18 ^(6a)	18 ^(6a)
Non-hazardous pollutant	Specific pollutant	Methiocarb	-	0.1 ⁽²⁾ (‘other individual pesticide’)	0.01 ^(6a)	-
Non-hazardous pollutant	Specific pollutant	Pendimethalin	-	0.1 ⁽²⁾ (‘other individual pesticide’)	0.3 ^(6a)	-

Substance classification		Determinant	Target concentrations (µg/l)			
Groundwater receptors ⁽⁵⁾	Surface water receptors ⁽⁶⁾		Minimum reporting value	UK drinking water standard (or best equivalent)	EQS or best equivalent	
					Freshwater	Transitional (estuaries) and coastal waters
Hazardous substance	Specific pollutant	Permethrin	0.001 ⁽⁷⁾	0.1 ⁽²⁾ (‘other individual pesticide’)	0.001 ^(6a)	0.0002 ^(6a)
Hazardous substance	Priority substance	Alachlor	-	0.1 ⁽²⁾ (‘other individual pesticide’)	0.3 ^(6a)	0.3 ^(6a)
Hazardous substance	Priority substance	Atrazine	0.03 ⁽⁷⁾	100 ⁽⁴⁾ (‘other individual pesticide’)	0.6 ^(6a)	0.6 ^(6a)
Hazardous substance	Priority substance	Diuron	-	0.1 ⁽²⁾ (‘other individual pesticide’)	0.2 ^(6a)	0.2 ^(6a)
Hazardous substance	Priority hazardous substance	Endosulphan	0.005 ⁽⁷⁾	0.1 ⁽²⁾ (‘other individual pesticide’)	0.005 ^(6a)	0.0005 ^(6a)
Non-hazardous pollutant	Priority substance	Isoproturon	-	0.1 ⁽²⁾ (‘other individual pesticide’)	0.3 ^(6a)	0.3 ^(6a)
Hazardous substance	Priority substance	Simazine	0.03 ⁽⁷⁾	0.1 ⁽²⁾ (‘other individual pesticide’)	1 ^(6a)	1 ^(6a)
Hazardous substance	Priority hazardous substance	Trifluralin	0.01 ⁽⁷⁾	0.1 ⁽²⁾ (‘other individual pesticide’)	0.03 ^(6a)	0.03 ^(6a)

Substance classification		Determinant	Target concentrations (µg/l)			
Groundwater receptors ⁽⁵⁾	Surface water receptors ⁽⁶⁾		Minimum reporting value	UK drinking water standard (or best equivalent)	EQS or best equivalent	
					Freshwater	Transitional (estuaries) and coastal waters
<i>(Not determined)</i>	From 22/12/18: Priority substance	Dichlorvos	-	0.1 ⁽²⁾ (‘other individual pesticide’)	From 22/12/18: 6.0E ^{-4(6a)}	From 22/12/18: 6.0E ^{-5(6a)}
Hazardous substance	From 22/12/18: Priority substance	Heptachlor and heptachlor epoxide	-	0.03 ⁽²⁾	From 22/12/18: 2.0E ^{-7(6a)}	From 22/12/18: 1.0E ^{-08(6a)}
Miscellaneous						
<i>None</i>	Specific pollutant	Triclosan (antibacterial agent)	-	-	0.1 ^(6a)	0.1 ^(6a)
Hazardous substance	From 22/12/18: Priority hazardous substance	Perfluoro-octane sulfonic acid (and its derivatives) (PFOS)	-	-	From 22/12/18: 6.5E ^{-4(6a)}	From 22/12/18: 1.3E ^{-4(6a)}
Hazardous substance	From 22/12/18: Priority hazardous substance	Hexabromo cyclododecane (HBCDD)	-	-	From 22/12/18: 0.0016 ^(6a)	From 22/12/18: 0.0008 ^(6a)

Substance classification		Determinant	Target concentrations (µg/l)			
Groundwater receptors ⁽⁵⁾	Surface water receptors ⁽⁶⁾		Minimum reporting value	UK drinking water standard (or best equivalent)	EQS or best equivalent	
					Freshwater	Transitional (estuaries) and coastal waters

Notes:

‘-’ A target concentration is not available.

\$Please note that total ammonia (NH₄⁺ and NH₃) is equivalent to ammoniacal nitrogen in laboratory reports

*¹ Please note that although iron is listed in the 2015 Direction as 1.000 µg/l, the EQS remains at 1mg/l in Scotland and it is assumed this is an error and should read either 1,000 or 1000µg/l.

*² Please note that although Isodrin is not listed in name within the group of “Cyclodiene pesticides” in Table 1 of Schedule 3 Part 3 of the 2015 Direction⁽⁶⁾, the CAS number for Isodrin (465-73-6) is listed and therefore it is assumed that it has been missed off the named list of substances.

*³ Total petroleum hydrocarbons is used for consistency, but is an analytical method-defined measurement for a mixture of hydrocarbons subject to environmental analysis¹¹.

“Bioavailable” in relation to copper, zinc, nickel and manganese (but not lead) is the generic EQSbioavailable^(6a) derived from the Metal Bioavailability Assessment Tool (M-BAT) developed by the Water Framework Directive UK Technical Advisory Group (WFDTAG). Exceedance of this value should prompt a site-specific assessment using the M-BAT with pH, DOC and Ca to derive a site-specific EQS termed the PNEC_{dissolved}.
<http://www.wfduk.org/resources/rivers-lakes-metal-bioavailability-assessment-tool-m-bat>.

For zinc, if there is an exceedance of the EQSbioavailable in an initial GQRA, Tier 2 required that the EQS for zinc should also have the ambient background concentration of zinc added as well

Table 2: World Health Organization (WHO) guide values for TPH CWG fractions in drinking water⁽¹³⁾ (as referenced in CL:AIRE, 2017⁽¹¹⁾)

TPH CWG fraction	WHO guide value for drinking water ⁽¹³⁾ (µg/l)
Aliphatic fractions:	
Aliphatic EC5-EC6	15,000
Aliphatic >EC6-EC8	15,000
Aliphatic >EC8-EC10	300
Aliphatic >EC10-EC12	300
Aliphatic >EC12-EC16	300
Aliphatic >EC16-EC21	-
Aliphatic >EC21-EC35	-
Aromatic fractions:	
Aromatic EC5-EC6	10 (benzene)
Aromatic >EC6-EC8	700 (toluene)
Aromatic >EC8-EC10	300 (ethyl benzene) 500 (xylenes)
Aromatic >EC10-EC12	90
Aromatic >EC12-EC16	90
Aromatic >EC16-EC21	90
Aromatic >EC21-EC35	90
Reference: World Health Organisation (WHO), 2008. Petroleum products in drinking-water. Background document for development of WHO guidelines for drinking water quality. WHO/SDE/WSH/05.08/123. World Health Organisation, Geneva ⁽¹³⁾ .	


References

1. Environment Agency (2017), 'The Environment Agency's approach to groundwater protection', version 1.0, March 2017 (formerly contained within GP3) [accessed 29 March 2017].
<https://www.gov.uk/government/collections/groundwater-protection>
- 1a. Environment Agency (2017), 'Land contamination groundwater compliance points: quantitative risk assessments', March 2017 (formerly contained within GP3) [accessed 29 March 2017].
<https://www.gov.uk/government/collections/groundwater-protection>
2. The Water Supply (Water Quality) Regulations 2016 (SI 2016/619)
 - 2a. Sum of chloroform, bromoform, dibromochloromethane and bromodichloromethane
 - 2b. Standard applies to individual pesticides except aldrin, dieldrin, heptachlor and heptachlor epoxide, for which a separate standard is defined.
3. The Private Water Supplies (England) Regulations 2016. SI 2016 / 618
4. WHO (2011), *Guidelines for drinking-water quality*, 4th edn
5. JAGDAG hazard substance determinations: This list contains substances that are determined to be hazardous substances or non-hazardous pollutants for the purposes of the groundwater directive 2006/118/EC. The absence of an assessment or substance from the list means an assessment has not been done yet and is presented as 'Not yet determined'; if a substance has been assessed but does not fall into either category it is presented as 'None'. For further details on how substances are assessed, see the Joint Agencies Groundwater Directive Advisory Group (JAGDAG) 'Methodology for the determination of hazardous substances in groundwater for the purposes of the groundwater directive 2006/118/EC' which is available from the JAGDAG website. The methodology is a UK –wide framework that sets criteria for how to assess whether a substance is a hazardous substances in groundwater. The list of substances can be found at:
<https://www.wfduk.org/stakeholders/jagdag>
6. The Water Framework Directive (Standards and Classification) Directions (England and Wales) 2015.
 - 6a. The EQS for these substances are based on a "long term mean" or an "annual average (AA)" EQS.
 - 6b. For cadmium and its compounds the EQS values vary depending on the hardness of the water as specified in five class categories (Class 1: < 40 mg CaCO₃/l, Class 2: 40 to < 50 mg CaCO₃/l, Class 3: 50 to < 100 mg CaCO₃/l, Class 4: 100 to < 200 mg CaCO₃/l and Class 5: ≥ 200 mg CaCO₃/l).
 - 6c. The EQS for Mercury and hexachlorobutadiene are based on a "maximum acceptable concentration (MAC)" EQS in absence of an "annual average (AA)" EQS.
 - 6d. The EQS for chlorine in saltwater is based on the 95th percentile concentration of total residual oxidant, which refers to the sum of all oxidising agents existing in water, expressed as available chlorine.
 - 6e. The recommended saltwater standard is derived using a safety factor of 100. Where the standard is failed, it is recommended that supporting evidence of ecological damage should be obtained before committing to expensive action.
 - 6f. EQS for total ammonia is as per Schedule 3, Part 1, Table 7 of of the above directions. EQS applies to river types 1, 2 and 4 and 6 (namely upland and low alkalinity). The EQS for a lowland and high alkalinity rivers (types 3, 5 and 7) is 600µg/l (0.6mg/l).

Additional information on the Metal Bioavailability Assessment Tool (M-BAT) is available at <http://www.wfduk.org/resources/rivers-lakes-metal-bioavailability-assessment-tool-m-bat>

7. Minimum reporting values listed at <https://www.gov.uk/government/publications/values-for-groundwater-risk-assessments/hazardous-substances-to-groundwater-minimum-reporting-values> (updated 13 January 2017; accessed 29 March 2017). Note target concentration for xylenes is 3 µg/l each for o-xylene and m/p xylene as it may not be possible to separate m- and p-xylene; 135 tcb, 124 tcb, 123 tcb each to 0.01 µg/l)
8. The Surface Waters (Abstraction for Drinking Water) (Classification) Regulations 1996 (as amended). SI 1996 / 3001
9. Council Directive on the Quality of Fresh Waters Needing Protection or Improvement in Order to Support Fish Life (Freshwater Fish Directive) (78/659/EEC)
10. WRc plc (2002), R&D Technical Report P45.
11. CL:AIRE, 2017. Petroleum Hydrocarbons in Groundwater: Guidance on assessing petroleum hydrocarbons using existing hydrogeological risk assessment methodologies. V1.1.
12. Drinking Water Inspectorate (London, UK). Environmental Information Request on MTBE in drinking water. Ref. DWI 1/10/18; dated 28 November 2006. Value is based on the odour threshold for MTBE, which is lower than a health-based guideline value
13. World Health Organisation (WHO), 2008. Petroleum products in drinking-water. Background document for development of WHO guidelines for drinking water quality. WHO/SDE/WSH/05.08/123. World Health Organisation, Geneva. [accessed 29 March 2017] http://www.who.int/water_sanitation_health/dwq/chemicals/petroleumproducts_2add_june2008.pdf

FLOW CHART TO ASSIST WITH SELECTION OF TARGET CONCENTRATIONS



TC = Target concentration

When leachate is being assessed the 'compliance point' is the groundwater body. Therefore dilution within the groundwater body may be applied with caution before comparing with the TC.

When directly assessing a receptor, e.g., a river, the appropriate TC should be selected.